

Dr. Duke's Phytochemical and Ethnobotanical Databases

List of Plants for T-CADINOL

Plant	Part	Low PPM	High PPM	StdDev	Reference
Satureja obovata	Leaf				Arrebola, M.L., Navaro, M.C., Jimenez, J. and Ocana, F.A. 1994. Variations in Yield and Composition of the Essential Oil of Satureja obovata. Phytochemistry 35(1): 83.
Satureja obovata	Leaf				Arrebola, M.L., Navaro, M.C., Jimenez, J. and Ocana, F.A. 1994. Variations in Yield and Composition of the Essential Oil of Satureja obovata. Phytochemistry 35(1): 83.
Satureja obovata	Leaf				Arrebola, M.L., Navaro, M.C., Jimenez, J. and Ocana, F.A. 1994. Variations in Yield and Composition of the Essential Oil of Satureja obovata. Phytochemistry 35(1): 83.
Vitex agnus-castus	Flower				Jim Duke's personal files.
Vitex agnus-castus	Fruit				Jim Duke's personal files.
Melia azedarach	Wood				--
Pinus sylvestris	Leaf				Leung, A. Y. and Foster, S. 1995. Encyclopedia of Common Natural Ingredients 2nd Ed. John Wiley & Sons, New York. 649 pp.
Hyssopus officinalis	Shoot				Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot				Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot				Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Cinnamomum camphora	Plant				--

Plant	Part	Low PPM	High PPM	StdDev	Reference
Satureja obovata	Leaf		0.01	-0.7680046689722265	Arrebola, M.L., Navaro, M.C., Jimenez, J. and Ocana, F.A. 1994. Variations in Yield and Composition of the Essential Oil of Satureja obovata. Phytochemistry 35(1): 83.
Ocimum basilicum	Plant		0.5	-0.7318552362623783	Die Nahrung. Pino, J., Rosado, A., Goire, I., Roncal, E., and Garcia, I. 1993. Analysis of the Essential Oil from Cuban Basil. Die Nahrung 37:(5): 501-504.
Ocimum basilicum	Plant		0.5	-0.7318552362623783	Die Nahrung. Pino, J., Rosado, A., Goire, I., Roncal, E., and Garcia, I. 1993. Analysis of the Essential Oil from Cuban Basil. Die Nahrung 37:(5): 501-504.
Teucrium salviastrum	Shoot		0.65	-1.0001295554196887	Velasco-Negueruela, A. and Perez-Alonso, M.J. 1990. The Volatiles of Six Teucrium Species from the Iberian Peninsula and the Balearic Islands. Phytochemistry 29(4): 1165-9.
Teucrium pseudoscorodonia	Shoot		0.82	-0.990356826743242	Velasco-Negueruela, A. and Perez-Alonso, M.J. 1990. The Volatiles of Six Teucrium Species from the Iberian Peninsula and the Balearic Islands. Phytochemistry 29(4): 1165-9.
Teucrium oxylepis	Shoot		1.0	-0.9800092316740633	Velasco-Negueruela, A. and Perez-Alonso, M.J. 1990. The Volatiles of Six Teucrium Species from the Iberian Peninsula and the Balearic Islands. Phytochemistry 29(4): 1165-9.
Teucrium scorodonia	Shoot		1.02	-0.978859498888599	Velasco-Negueruela, A. and Perez-Alonso, M.J. 1990. The Volatiles of Six Teucrium Species from the Iberian Peninsula and the Balearic Islands. Phytochemistry 29(4): 1165-9.
Teucrium asiaticum	Shoot		1.16	-0.9708113693903488	Velasco-Negueruela, A. and Perez-Alonso, M.J. 1990. The Volatiles of Six Teucrium Species from the Iberian Peninsula and the Balearic Islands. Phytochemistry 29(4): 1165-9.
Salvia dorisiana	Shoot	1.2	1.8	-0.9340199202554911	Tucker, A.O. & Maciarelo, M.J. 1994. The Essential Oil of Salvia dorisiana Standley. J. Ess. Oil Res. 6: 97-8.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Vitex agnus-castus	Leaf		2.5	-0.7272372511202277	Ekundayo, O., Laakso, I., Holopainen, M., Hiltunen, R., Oguntimein, B., and Kauppinen, V. 1990. The Chemical Composition and Antimicrobial Activity of the Leaf Oil of <i>Vitex agnus-castus</i> L. J. Essential Oil Research, 2: 115-119.
Teucrium gnaphalodes	Shoot		2.5	-0.8937792727642405	Perez-Alonso, M.J. Velasco-Negueruela, A. and Lopez-Saez, J.A. 1993. The Essential Oils of Two Iberian Teucrium Species. J. Ess. Oil Res. 5: 397-402.
Acinos alpinus	Shoot		2.5	-0.8937792727642405	Velasco-Negueruela, A., Perez-Alonso, M.J., Jimenez, S.M. and Garcia, F.M. 1993. The Volatile Constituents of <i>Acinus alpinus</i> (L.) Moench ssp. meridionalis (Nyman). P.W. Ball Growing in Spain. Flav. & Frag. J. 8:127-130.)
Acinos alpinus	Shoot		2.5	-0.8937792727642405	Velasco-Negueruela, A., Perez-Alonso, M.J., Jimenez, S.M. and Garcia, F.M. 1993. The Volatile Constituents of <i>Acinus alpinus</i> (L.) Moench ssp. meridionalis (Nyman). P.W. Ball Growing in Spain. Flav. & Frag. J. 8:127-130.)
Teucrium polium	Shoot		3.0	-0.8650359531276329	Perez-Alonso, M.J. Velasco-Negueruela, A. and Lopez-Saez, J.A. 1993. The Essential Oils of Two Iberian Teucrium Species. J. Ess. Oil Res. 5: 397-402.
Tagetes lucida	Shoot		6.0	-0.6925760353079873	Bicchi, C., Fresia, M., Rubiolo, P., Monti, D., Franz, C., Goehler, I. 1997. Constituents of <i>Tagetes lucida</i> Cav. ssp. <i>lucida</i> essential oil. Flavor & Fragrance, 12(1): 47-52.
Humulus lupulus	Fruit		9.0		--
Teucrium oxylepis	Shoot		9.24	-0.5063193240627701	Velasco-Negueruela, A. and Perez-Alonso, M.J. 1990. The Volatiles of Six Teucrium Species from the Iberian Peninsula and the Balearic Islands. Phytochemistry 29(4): 1165-9.
Achillea millefolium	Leaf	1.0	15.0	-0.5225815390117998	--
Hyssopus officinalis	Shoot		20.0	0.1122369145170252	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (<i>Hyssopus officinalis</i> L.) J. Agric. Food Chem. 42: 776-781.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Hyssopus officinalis	Shoot		20.0	0.1122369145170252	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Melissa officinalis	Shoot	3.0	24.0	0.342183471609886	Deutsche Apot. Ziti. 129(4):155-163. W. Schulze et al. Die Melisse.
Thymus funkii	Shoot		25.0	0.39967011088310117	Vila, R., et al. 1995. Composition and study of the variability of the essential oil of Thymus funkii Cousson. Flav. & Fragr. J. 10(6): 379-383.
Thymus riatarum	Shoot		25.0	0.39967011088310117	Iglesias, J., Vila, R., Canigueral, S., Bellakdhar, and Il Idrissi, A. 1991. Analysis of the Essential Oil of Thymus riatarum. J. Ess. Oil Res. 3: 43-4.
Stevia rebaudiana	Leaf		28.0	-0.309739598419035	Kinghorn, A. D. (Ed.) 2002. Medicinal and Aromatic Plants - Industrial Profiles. Stevia. The genus Stevia. Taylor & Francis. New York, NY. 211 pp.
Hyssopus officinalis	Shoot		30.0	0.6871033072491771	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot		30.0	0.6871033072491771	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot		30.0	0.6871033072491771	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot		30.0	0.6871033072491771	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot		30.0	0.6871033072491771	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Thymus funkii	Shoot		35.0	0.9745365036152531	Vila, R., et al. 1995. Composition and study of the variability of the essential oil of Thymus funkii Cousson. Flav. & Fragr. J. 10(6): 379-383.
Aloysia citrodora	Plant	5.0	35.0	-0.23187492634055548	Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
Nepeta racemosa	Shoot		60.0	2.411702485445633	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.
Nepeta racemosa	Shoot		60.0	2.411702485445633	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.
Stevia rebaudiana	Flower		60.0		Kinghorn, A. D. (Ed.) 2002. Medicinal and Aromatic Plants - Industrial Profiles. Stevia. The genus Stevia. Taylor & Francis. New York, NY. 211 pp.
Hyssopus officinalis	Leaf	0.1	60.0	0.2141790245785401	Flavour and Fragrance Journal, 6: 72.
Artemisia pallens	Plant	38.0	168.0	1.6955853988653122	--
Metrosideros nervulosa	Leaf		176.0	2.1133840329447495	--
Pinus sylvestris	Leaf Essent. Oil		2000.0	-1.0	--
Vitex agnus-castus	Leaf Essent. Oil		10700.0	1.0	Jim Duke's personal files.
Plectranthus barbatus	Essential Oil	5000.0	41000.0		--
Ocimum basilicum	Shoot Essent. Oil		71400.0		--