

Dr. Duke's Phytochemical and Ethnobotanical Databases

List of Plants for TRICYCLEN

Plant	Part	Low PPM	High PPM	StdDev	Reference
Abies alba	Leaf	45.0	63.0	0.5496290114333532	--
Cymbopogon nardus	Plant	45.0	180.0		Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
Salvia officinalis	Leaf	15.0	84.0	1.3596086072298736	Duke, J. A. Writeups or information summaries on approximately 2,000 economic plants, USDA, ARS, Beltsville, MD 20705.
Origanum minutiflorum	Shoot	5.0	7.0	-0.25899623151650575	Baser, K.H.C., Tumen, G., Sezik, E. 1991. The Essential Oil of Origanum minutiflorum O. Schwarz and P.H. Davis. J. Ess. Oil Res. 3: 445-446.
Zingiber officinale	Rhizome	2.0	115.0		--
Achillea millefolium	Leaf	0.6	27.0	-0.8389074385035388	Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
Carum carvi	Seed				Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
Ocimum basilicum	Essential Oil				--
Rosmarinus officinalis	Leaf Essent. Oil				--
Zingiber officinale	Rhizome Essent. Oil		2300.0		--
Lavandula x intermedia	Plant				Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
Thymus longicaulis	Shoot		0.0	-0.2829643095555945	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot		1.7	-0.2771434906032444	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot		0.0	-0.2829643095555945	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Rosmarinus officinalis	Plant				Duke, J. A. Writeups or information summaries on approximately 2,000 economic plants, USDA, ARS, Beltsville, MD 20705.
Origanum syriacum	Shoot		0.0	-0.2829643095555945	Fleisher, A. & Fleisher, Z. 1991. Chemical Composition of Origanum syriacum L. Essential Oil. J. Ess. Oil Res. 3: 121-123.
Origanum syriacum	Shoot		3.0	-0.27269227611027075	Fleisher, A. & Fleisher, Z. 1991. Chemical Composition of Origanum syriacum L. Essential Oil. J. Ess. Oil Res. 3: 121-123.
Origanum syriacum	Shoot		0.0	-0.2829643095555945	Fleisher, A. & Fleisher, Z. 1991. Chemical Composition of Origanum syriacum L. Essential Oil. J. Ess. Oil Res. 3: 121-123.
Carum carvi	Fruit				--
Pinus sylvestris	Essential Oil				--
Pinus sylvestris	Leaf Essent. Oil		4400.0	1.0	--
Pinus sylvestris	Resin, Exudate, Sap				--
Salvia canariensis	Leaf		21.0	-1.0703301801596874	Casnigueral, S., Iglesias, J., Vila, R., Virgili, A. and Ibanez, C. 1994. The Essential Oil from Leaves of Salvia canariensis L. Flav. & Fragr. J. 9:201-204. S. Canigueral, Facultat de Farmacia, Universitat de Barcelona, Ave. Diagonal 643, E-08028, Barcelona Spain
Crocus sativus	Silk Stigma Style				--
Ocimum gratissimum	Shoot		0.1	-0.2826219084407503	Vostrowsky, O., Garbe, W., Bestmann, H.J. and Maia, J.G.S. 1990. Essential Oil of Alfavaca, Ocimum gratissimum, from Brazilian Amazon. Zeitschr. Naturforschung 45(C): 1073-6.
Thymus funkii	Shoot		24.0	-0.20078804199300457	Vila, R., et al. 1995. Composition and study of the variability of the essential oil of Thymus funkii Cousson. Flav. & Fragr. J. 10(6): 379-383.
Thymus funkii	Shoot		24.0	-0.20078804199300457	Vila, R., et al. 1995. Composition and study of the variability of the essential oil of Thymus funkii Cousson. Flav. & Fragr. J. 10(6): 379-383.
Juniperus communis	Fruit				--

Plant	Part	Low PPM	High PPM	StdDev	Reference
Ferula gummosa	Plant				--
Thymus riatarum	Shoot		0.1	-0.2826219084407503	Iglesias, J., Vila, R., Canigueral, S., Bellakdhar, and I Idrissi, A. 1991. Analysis of the Essential Oil of Thymus riatarum. J. Ess. Oil Res. 3: 43-4.
Origanum sipyleum	Shoot		30.0	-0.1802439751023571	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum sipyleum	Shoot		20.0	-0.21448408658676957	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum onites	Shoot		15.0	-0.2316041423289758	Biondi, D., Cianci, P., Geraci, C. and Ruberto, G. 1993. Antimicrobial Activity and Chemical Composition of Essential Oils from Sicilian Aromatic Plants. Flav. & Frag. J. 8: 331-7.
Origanum sipyleum	Shoot		30.0	-0.1802439751023571	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum sipyleum	Shoot				Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Thymus capitatus	Shoot		0.0	-0.2829643095555945	Biondi, D., Cianci, P., Geraci, C. and Ruberto, G. 1993. Antimicrobial Activity and Chemical Composition of Essential Oils from Sicilian Aromatic Plants. Flav. & Frag. J. 8: 331-7.
Salvia officinalis	Leaf Essent. Oil		3000.0	-1.0	--
Picea mariana	Twig		46.0		--
Artemisia salsoloides	Shoot		1250.0	3.997049625995964	V. Kaul, P. Weyerstahl, H. Wahlberg, H. Marschall, (1992); Volatile constituents of the essential oil and the absolute of Artemisia salsoloides Willd. from Ladakh, Flavour and Fragrance journal, Vol.7, 299-305.