

Dr. Duke's Phytochemical and Ethnobotanical Databases

List of Plants for (Z)-BETA-OCIMENE

Plant	Part	Low PPM	High PPM	StdDev	Reference
Agastache foeniculum	Plant	3.0	120.0	1.4074844462950056	--
Alpinia galanga	Rhizome	8.0	985.0		--
Alpinia galanga	Leaf		1705.0	1.4142018319896454	--
Apium graveolens	Seed	6.0	9.0		Charalambous, G. (Ed.). 1994. Spices, Herbs and Edible Fungi. Elsevier Science B. V. Amsterdam. 764 pp.
Calamintha nepeta	Shoot		9.0	-0.40626642590156903	Kirimer, N., Baser, K.H.C., Ozek, T. and Kurkcuoglu, M. 1992. Composition of the Essential Oil of Calamintha nepeta subsp. glandulosa. J. Ess. Oil Res. 4:189-190
Carica papaya	Fruit				--
Carum carvi	Seed				--
Cinnamomum verum	Leaf		0.0	-0.7120892822452273	Mallavarupu, G. R. et al. 1995. Investigation of the essential oil of cinnamon leaf grown at Bangalore and Hyderabad. Flav. & Fragr. J., 10: 239-242.
Citrus paradisi	Pericarp		1.0		--
Citrus sinensis	Fruit	2.0	3.0	-0.6648405774204937	--
Citrus limon	Petiole		3.5		--
Citrus reticulata	Fruit		0.1	-0.7485469887599769	--
Dictamnus albus	Shoot		15.0	-0.37869400950362786	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Planta Med. 60:481-2
Dictamnus albus	Shoot		0.5	-0.4453273491319857	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Planta Med. 60:481-2
Dictamnus albus	Shoot		15.0	-0.37869400950362786	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Planta Med. 60:481-2
Elsholtzia blanda	Shoot		18.0	-0.3649078013046573	Bestman, H.J., Rauscher, J., Vostrowsky O., Pant, A.K., Dev. V., Perihar, R. and Mathela, C.S. 1992. Constituents of the Essential Oil of Elsholtzia blanda Benth. (Labiatae). J. Ess. Oils Res. 4: 121-124

Plant	Part	Low PPM	High PPM	StdDev	Reference
Hyptis suaveolens	Shoot		3.0	-0.4338388422995102	Mallvarapu, G.R., Ramesh, S., Kaul, P.N., Bhattacharya, A.K., and Rao, B.R.R. 1993. The Essential Oil of Hyptis suaveolens (L.) Poit. J. Ess. Oil Res. 5: 321.
Hyptis suaveolens	Shoot		3.0	-0.4338388422995102	Mallvarapu, G.R., Ramesh, S., Kaul, P.N., Bhattacharya, A.K., and Rao, B.R.R. 1993. The Essential Oil of Hyptis suaveolens (L.) Poit. J. Ess. Oil Res. 5: 321.
Hyssopus officinalis	Shoot		980.0	4.055869627831911	Tsankova, E.T., Konatchiev, A.N. and Genova, E.M. 1993. Chemical Composition of the Essential Oils of Two Hyssopus officinalis cultivars. J. Ess. Oil Res. 5: 609-611.
Hyssopus officinalis	Shoot		690.0	2.7232028352647544	Tsankova, E.T., Konatchiev, A.N. and Genova, E.M. 1993. Chemical Composition of the Essential Oils of Two Hyssopus officinalis cultivars. J. Ess. Oil Res. 5: 609-611.
Leonotis leonurus	Se		24.0		Pedro, L.G., Barroso, J.G., Marques, N.T., Ascensao, L., Pais, M.S.S. and Scheffer, J.J.C. 1991. Composition of the Essential Oil from Sepals of Leonotis leonurus R. Br. J. Ess. Oil Res. 3: 451-3
Lindera benzoin	Fruit	55.0	75.0	1.413387566180471	--
Minthostachys mollis	Shoot	4.0	8.0	-0.4108618286345592	Alkire, B.H., Tucker, A.O., and Maciarelo, M.J. 1994. Tipo (Minthostachys mollis (Lamiaceae): An Ecuadorian Mint. Econ. Bot. 48(1): 60-64.
Nepeta racemosa	Shoot		33.0	-0.29597676030980435	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.
Nepeta racemosa	Shoot		33.0	-0.29597676030980435	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.
Origanum vulgare	Plant		5.0	-0.7847599508687727	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Origanum vulgare	Plant		20.0	-0.4988150294995841	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum sipyleum	Shoot		15.0	-0.37869400950362786	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum vulgare	Plant		5.0	-0.7847599508687727	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum sipyleum	Shoot		7.0	-0.4154572313675494	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum sipyleum	Shoot		205.0	0.49443250976450925	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum vulgare	Plant		7.0	-0.7466339613528808	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum sipyleum	Shoot		205.0	0.49443250976450925	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Pelargonium citrosum	Shoot		3.0	-0.4338388422995102	Matsuda, B. M., et al. 1996. Essential Oil Analysis and Field Evaluation of the Citrosa Plant 'Pelargonium citrosum' as a Repellent Against Populations of Aedes Mosquitoes. J. Am. Mosq. Contr. Assoc. 12(1):69-74.
Porophyllum ruderale	Plant	80.0	120.0	1.4074844462950056	Loayza, I., de Groot, W., Lorenzo, D. et al. 1999. Composition of the essential oil of Porophyllum ruderale (Jacq.) Cass. from Bolivia. Flav. & Fragr. J. 14: 393-8.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Rosmarinus officinalis	Shoot		13.0	-0.38788481496960825	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus x lavandulaceus	Shoot	26.0	110.0	0.05786925013044062	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus eriocalyx	Shoot				Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus tomentosus	Shoot		0.1	-0.4471655102251818	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot				Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus eriocalyx	Shoot		0.1	-0.4471655102251818	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot	6.0	75.0	-0.10296984552421623	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.

Plant	Part	Low PPM	High PPM	StdDev	Reference
<i>Rosmarinus x mendizabalii</i>	Shoot		0.1	-0.4471655102251818	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the <i>Rosmarinus eriocalyx-tomentosus</i> Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
<i>Rosmarinus officinalis</i>	Shoot	6.0	75.0	-0.10296984552421623	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the <i>Rosmarinus eriocalyx-tomentosus</i> Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
<i>Sassafras albidum</i>	Leaf	2.6	8.0	-0.7021125497444185	--
<i>Satureja cilicica</i>	Shoot		210.0	0.5174095234294602	Tumen, G. Baser, K.H.C. and Kirimer, N. 1993. The Essential Oil of <i>Satureja cilicica</i> P.H. Davis. J. Ess. Oil Res. 5: 547-548.
<i>Tagetes lucida</i>	Shoot				Bicchi, C., Fresia, M., Rubiolo, P., Monti, D., Franz, C., Goehler, I. 1997. Constituents of <i>Tagetes lucida</i> Cav. ssp. <i>lucida</i> essential oil. Flavor & Fragrance, 12(1): 47-52.
<i>Thymus longicaulis</i>	Shoot		0.8	-0.4439487283120886	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of <i>Thymus longicaulis</i> C. Presl subsp. <i>longicaulis</i> in the same Population. J. Ess. Oil Res. 5: 291-5.
<i>Thymus longicaulis</i>	Shoot		0.8	-0.4439487283120886	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of <i>Thymus longicaulis</i> C. Presl subsp. <i>longicaulis</i> in the same Population. J. Ess. Oil Res. 5: 291-5.
<i>Thymus longicaulis</i>	Shoot		0.0	-0.4476250504984808	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of <i>Thymus longicaulis</i> C. Presl subsp. <i>longicaulis</i> in the same Population. J. Ess. Oil Res. 5: 291-5.