

Dr. Duke's Phytochemical and Ethnobotanical Databases

List of Plants for 2,4-DECADIENAL

Plant	Part	Low PPM	High PPM	StdDev	Reference
Lonicera japonica	Flower		0.001		Schlotzhauer, W.S., S.D. Pair, and R.J. Horvat. 1996. Volatile constituents from the flowers of Japanese Honeysuckle. J. Agric. Food Chem. 44:206-209.
Mentha aquatica	Shoot				Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
Mentha aquatica	Shoot		2.0	1.0000000000000002	Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
Mentha aquatica	Shoot				Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
Mentha aquatica	Shoot				Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
Mentha aquatica	Shoot		0.1	-1.0	Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
Vanilla planifolia	Fruit		25.0		--