

Dr. Duke's Phytochemical and Ethnobotanical Databases

List of Plants for CIS-SABINENE-HYDRATE

Plant	Part	Low PPM	High PPM	StdDev	Reference
Origanum majorana	Leaf Essent. Oil	20000.0	580000.0		--
Origanum majorana	Essential Oil		75000.0	1.0	--
Rosmarinus officinalis	Shoot Essent. Oil		19300.0	1.0	--
Origanum majorana	Leaf	100.0	7965.0	3.162122330899287	--
Myristica fragrans	Seed Essent. Oil	2000.0	7000.0	1.0	--
Myristica fragrans	Essential Oil	2000.0	5000.0	-1.0	--
Origanum vulgare	Shoot Essent. Oil		2200.0	-1.0	--
Melaleuca alternifolia	Root Essent. Oil		2000.0		--
Artemisia salsoloides	Shoot		2000.0	5.278612667288771	V. Kaul, P. Weyerstahl, H. Wahlberg, H. Marschall, (1992); Volatile constituents of the essential oil and the absolute of <i>Aremisia salsoloides</i> Willd. from Ladakh, Flavour and Fragrance journal, Vol.7, 299-305.
Cuminum cyminum	Seed Essent. Oil		1500.0	-1.0	--
Monarda fistulosa	Plant	1.0	527.0	2.6281938059149086	Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
Cuminum cyminum	Seed		207.0		Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
Cuminum cyminum	Fruit		207.0	1.414213562373095	--
Artemisia annua	Plant	1.0	200.0	0.6462396290052647	--
Achillea millefolium	Leaf	1.0	80.0	-0.2862032724587523	--
Origanum minutiflorum	Shoot	60.0	80.0	-0.027368494787144388	Baser, K.H.C., Tumen, G., Sezik, E. 1991. The Essential Oil of <i>Origanum minutiflorum</i> O. Schwarz and P.H. Davis. J. Ess. Oil Res. 3: 445-446.
Thymus funkii	Shoot		65.0	-0.06882147261586248	Vila, R., et al. 1995. Composition and study of the variability of the essential oil of <i>Thymus funkii</i> Cousson. Flav. & Fragr. J. 10(6): 379-383.
Thymus funkii	Shoot		65.0	-0.06882147261586248	Vila, R., et al. 1995. Composition and study of the variability of the essential oil of <i>Thymus funkii</i> Cousson. Flav. & Fragr. J. 10(6): 379-383.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Thymus cilicicus	Shoot		64.0	-0.07158500447111035	Tumen, G., Koyuncu, M., Kirimer, N., and Baser, K.H.C. 1994. Composition of the Essential Oil of Thymus cilicicus Boiss. & Bal. J. Ess. Oil Res. 6: 97-8.
Origanum vulgare	Plant		60.0	-0.2023034436777631	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum syriacum	Shoot		55.0	-0.0964567911683412	Fleisher, A. & Fleisher, Z. 1991. Chemical Composition of Origanum syriacum L. Essential Oil. J. Ess. Oil Res. 3: 121-123.
Origanum sipyleum	Shoot		50.0	-0.11027445044458056	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum sipyleum	Shoot		50.0	-0.11027445044458056	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum vulgare	Plant	1.0	46.0	-0.2871577509460659	Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
Origanum sipyleum	Shoot		45.0	-0.12409210972081992	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum sipyleum	Shoot		35.0	-0.15172742827329866	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Satureja cilicica	Shoot		21.0	-0.19041687424676887	Tumen, G. Baser, K.H.C. and Kirimer, N. 1993. The Essential Oil of Satureja cilicica P.H. Davis. J. Ess. Oil Res. 5: 547-548.
Micromeria teneriffae	Leaf		20.0	-0.31244290926299867	Kirimer, N., Ozek, T., and Baser, K.H.C. 1991. Composition of the Essential Oil of Micromeria congesta. J. Ess. Oil Res., 3: 387-393.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Thymus zygis	Shoot		20.0	-0.19318040610201673	Tantaoui-Elaraki, A., Lattaoui, N., Errifi, A. and Benjlali, B. 1993. Composition and Antimicrobial Activity of the Essential Oils of Thymus broussonettii, T. zygis and T. saturejoides. J. Ess. Oil Res. 5: 45-53.
Micromeria varia	Shoot		15.0	-0.2069980653782561	Pedro, L.G., et al. 1995. Composition of the Essential oil of Micromeria varia Benth. ssp. thymoides (Sol. ex Lowe) Perez var. thymoides, and endemic species of the Madeira Archipelago. flav. & Fragr. J. 10(3): 199-202.
Micromeria varia	Shoot		15.0	-0.2069980653782561	--
Murraya koenigii	Leaf		12.0	-0.31594152750356486	--
Thymus longicaulis	Shoot		7.0	-0.22910632022023908	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Micromeria fruticosa	Shoot		5.0	-0.23463338393073482	Fleisher, Z. and Fleisher, A. 1991. The Essential Oil of Micromeria fruticosa (L.) Druce subsp. barbata (Boiss et. Ky.), P.H. Davis. Aromatic Plants of the Holy Land and the Sinai. Part VII. J. Ess. Oil Res 3: 477-479.
Micromeria fruticosa	Shoot		5.0	-0.23463338393073482	Fleisher, Z. and Fleisher, A. 1991. The Essential Oil of Micromeria fruticosa (L.) Druce subsp. barbata (Boiss et. Ky.), P.H. Davis. Aromatic Plants of the Holy Land and the Sinai. Part VII. J. Ess. Oil Res 3: 477-479.
Calamintha nepeta	Shoot		4.0	-0.2373969157859827	Kirimer, N., Baser, K.H.C., Ozek, T. and Kurkcuoglu, M. 1992. Composition of the Essential Oil of Calamintha nepeta subsp. glandulosa. J. Ess. Oil Res. 4:189-190
Origanum onites	Plant		4.0	-0.5417206727509742	Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
Nepeta racemosa	Shoot		3.0	-0.24016044764123057	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.

Plant	Part	Low PPM	High PPM	StdDev	Reference
<i>Nepeta racemosa</i>	Shoot		3.0	-0.24016044764123057	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of <i>Nepeta racemosa</i> Lam. J. Ess. Oil Res. 5: 215-7.
<i>Sideritis germanicolpitana</i>	Plant	1.0	2.0	-0.5538427166464461	J. Essential Oil, 4: 533.
<i>Calamintha nepeta</i>	Leaf		1.0	-0.32075212758434335	Akgul, A., De Pooter, H.L., and De Buyck, L.F. 1991. The Essential Oils of <i>Calamintha nepeta</i> subsp. <i>glandulosa</i> and <i>Ziziphora clinopodioides</i> from Turkey. J. Ess. Oil Res., 3: 7-10.
<i>Origanum vulgare</i>	Plant		0.9	-0.5605098407889556	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four <i>Origanum vulgare</i> Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
<i>Origanum vulgare</i>	Plant		0.5	-0.5629342495680499	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four <i>Origanum vulgare</i> Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
<i>Rosmarinus officinalis</i>	Leaf		0.4	-0.3210145239523858	Soliman, F.M., El-Kashoury, M.M., Fathy, M.M. and Gonaid, M.H. 1994. Analysis and Biological Activity of the Essential Oil of <i>Rosmarinus officinalis</i> L. from Egypt. Flavour and Fragrance J. 9: 29-33.
<i>Rosmarinus officinalis</i>	Leaf		0.4	-0.3210145239523858	Soliman, F.M., El-Kashoury, M.M., Fathy, M.M. and Gonaid, M.H. 1994. Analysis and Biological Activity of the Essential Oil of <i>Rosmarinus officinalis</i> L. from Egypt. Flavour and Fragrance J. 9: 29-33.
<i>Citrus sinensis</i>	Fruit		0.1	-0.7071067811865477	--
<i>Micromeria myrtifolia</i>	Shoot		0.1	-0.24817469002144935	Ozek, T., Kirimer, N., and Baser, K.H.C. 1992. Composition of the Essential Oil of <i>Micromeria myrtifolia</i> Boiss. et Hohen. J. Ess. Oil Res., 4: 79-80.
<i>Thymus saturejoides</i>	Shoot		0.1	-0.24817469002144935	Tantaoui-Elaraki, A., Lattaoui, N., Errifi, A. and Benjilali, B. 1993. Composition and Antimicrobial Activity of the Essential Oils of <i>Thymus broussonettii</i> , <i>T. zygis</i> and <i>T. saturejoides</i> . J. Ess. Oil Res. 5: 45-53.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Citrus reticulata	Fruit		0.1	-0.7071067811865477	--
Satureja obovata	Leaf		0.01	-0.3211850815916133	Arrebola, M.L., Navaro, M.C., Jimenez, J. and Ocana, F.A. 1994. Variations in Yield and Composition of the Essential Oil of Satureja obovata. Phytochemistry 35(1): 83.
Elsholtzia polystachya	Leaf		0.0	-0.32118945486441414	Mathela, C.S., Melkani, A.B., Bisht, J.C., Pant, A.K., Bestmann, H.J., Erler, J., Kobold, U., Rauscher, J. and Vostrowsky, O. 1992. Chemical Varieties of Essential Oils from Elsholtzia polystachya from Two Different Locations in India. Planta Medica 58: 376-379.
Elsholtzia polystachya	Leaf		0.0	-0.32118945486441414	Mathela, C.S., Melkani, A.B., Bisht, J.C., Pant, A.K., Bestmann, H.J., Erler, J., Kobold, U., Rauscher, J. and Vostrowsky, O. 1992. Chemical Varieties of Essential Oils from Elsholtzia polystachya from Two Different Locations in India. Planta Medica 58: 376-379.
Origanum vulgare	Plant		0.0	-0.5659647605419179	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Thymus broussonettii	Shoot		0.0	-0.2484510432069742	Tantaoui-Elaraki, A., Lattaoui, N., Errifi, A. and Benjlali, B. 1993. Composition and Antimicrobial Activity of the Essential Oils of Thymus broussonettii, T. zygis and T. saturejoides. J. Ess. Oil Res. 5: 45-53.
Rosmarinus officinalis	Leaf		0.0	-0.32118945486441414	Soliman, F.M., El-Kashoury, M.M., Fathy, M.M. and Gonaid, M.H. 1994. Analysis and Biological Activity of the Essential Oil of Rosmarinus officinalis L. from Egypt. Flavour and Fragrance J. 9: 29-33.
Origanum onites	Shoot		0.0	-0.2484510432069742	Biondi, D., Cianci, P., Geraci, C. and Ruberto, G. 1993. Antimicrobial Activity and Chemical Composition of Essential Oils from Sicilian Aromatic Plants. Flav. & Frag. J. 8: 331-7.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Thymus capitatus	Shoot		0.0	-0.2484510432069742	Biondi, D., Cianci, P., Geraci, C. and Ruberto, G. 1993. Antimicrobial Activity and Chemical Composition of Essential Oils from Sicilian Aromatic Plants. Flav. & Frag. J. 8: 331-7.
Thymus longicaulis	Shoot		0.0	-0.2484510432069742	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot		0.0	-0.2484510432069742	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Origanum syriacum	Shoot		0.0	-0.2484510432069742	Fleisher, A. & Fleisher, Z. 1991. Chemical Composition of Origanum syriacum L. Essential Oil. J. Ess. Oil Res. 3: 121-123.
Origanum syriacum	Shoot		0.0	-0.2484510432069742	Fleisher, A. & Fleisher, Z. 1991. Chemical Composition of Origanum syriacum L. Essential Oil. J. Ess. Oil Res. 3: 121-123.
Rosmarinus officinalis	Leaf Essent. Oil				--
Mentha spicata	Essential Oil				--
Angelica archangelica	Root Essent. Oil				--
Rosmarinus officinalis	Plant				--
Salvia officinalis	Leaf Essent. Oil				--
Piper nigrum	Fruit				--
Ocimum basilicum	Plant				--
Angelica archangelica	Root				Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
Satureja obovata	Leaf				Arrebola, M.L., Navaro, M.C., Jimenez, J. and Ocana, F.A. 1994. Variations in Yield and Composition of the Essential Oil of Satureja obovata. Phytochemistry 35(1): 83.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Satureja obovata	Leaf				Arrebola, M.L., Navaro, M.C., Jimenez, J. and Ocana, F.A. 1994. Variations in Yield and Composition of the Essential Oil of Satureja obovata. Phytochemistry 35(1): 83.
Satureja obovata	Leaf				Arrebola, M.L., Navaro, M.C., Jimenez, J. and Ocana, F.A. 1994. Variations in Yield and Composition of the Essential Oil of Satureja obovata. Phytochemistry 35(1): 83.
Laurus nobilis	Leaf				--
Myristica fragrans	Seed				Leung, A.Y., Encyclopedia of Common Natural Ingredients Used in Food, Drugs, and Cosmetics, John Wiley & Sons, New York, 1980.
Salvia officinalis	Leaf				Duke, J. A. Writeups or information summaries on approximately 2,000 economic plants, USDA, ARS, Beltsville, MD 20705.
Aralia cordata	Root				--