

Dr. Duke's Phytochemical and Ethnobotanical Databases

List of Plants for (E)-BETA-OCIMENE

Plant	Part	Low PPM	High PPM	StdDev	Reference
Ocimum basilicum	Plant	1.0	435.0	2.2262230322078222	--
Rosmarinus officinalis	Shoot		380.0	4.150712157567762	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus eriocalyx	Shoot	60.0	185.0	1.6725587938311206	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus eriocalyx	Shoot	60.0	185.0	1.6725587938311206	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Tagetes lucida	Shoot		175.0	1.5454740059471903	Bicchi, C., Fresia, M., Rubiolo, P., Monti, D., Franz, C., Goehler, I. 1997. Constituents of Tagetes lucida Cav. ssp. lucida essential oil. Flavor & Fragrance, 12(1): 47-52.
Thymus zygis	Shoot		75.0	0.27462612710788703	Jimenez, J., Navarro, M.C., Montilla, M.P., Martin, A. and Martinez, A. 1993. Thymus zygis Oil: Its Effects on CCl4-Induced Hepatotoxicity and Free Radical Scavenger Activity. JEO5: 153-8.
Origanum sipyleum	Shoot		70.0	0.21108373316592186	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum sipyleum	Shoot		70.0	0.21108373316592186	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Hyssopus officinalis	Shoot		70.0	0.21108373316592186	Tsankova, E.T., Konatchiev, A.N. and Genova, E.M. 1993. Chemical Composition of the Essential Oils of Two Hyssopus officinalis cultivars. J. Ess. Oil Res. 5: 609-611.

Plant	Part	Low PPM	High PPM	StdDev	Reference
<i>Hyssopus officinalis</i>	Shoot		70.0	0.21108373316592186	Tsankova, E.T., Konatchiev, A.N. and Genova, E.M. 1993. Chemical Composition of the Essential Oils of Two <i>Hyssopus officinalis</i> cultivars. <i>J. Ess. Oil Res.</i> 5: 609-611.
<i>Satureja cilicica</i>	Shoot		55.0	0.020456551340026382	Tumen, G. Baser, K.H.C. and Kirimer, N. 1993. The Essential Oil of <i>Satureja cilicica</i> P.H. Davis. <i>J. Ess. Oil Res.</i> 5: 547-548.
<i>Lindera benzoin</i>	Fruit	36.0	50.0	1.410387910020219	--
<i>Origanum vulgare</i>	Plant		50.0	-0.3058179350004837	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four <i>Origanum vulgare</i> Subspecies of Anatolian Origin. <i>J. Ess. Oil Res.</i> , 5: 425-431.
<i>Lavandula x hybrida</i>	Shoot	22.0	48.0	-0.06850280017872502	Tucker, A.O., Maciarelo, M.J., Angell, S., Espaillat, J.R., and French, E.C. 1993. The Essential Oil of <i>Lavandula x hybrida</i> Balb. ex Ging., a Distinct Hybrid from <i>L. x heterophylla</i> Poir. (Labiatae). <i>J. Ess. Oil Res.</i> 5: 443-445.
<i>Origanum vulgare</i>	Plant		40.0	-0.3715852328500501	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four <i>Origanum vulgare</i> Subspecies of Anatolian Origin. <i>J. Ess. Oil Res.</i> , 5: 425-431.
<i>Murraya koenigii</i>	Leaf		40.0	1.504389132845184	--
<i>Elsholtzia blanda</i>	Shoot		38.0	-0.19558758806265536	Bestman, H.J., Rauscher, J., Vostrowsky O., Pant, A.K., Dev. V., Perihar, R. and Mathela, C.S. 1992. Constituents of the Essential Oil of <i>Elsholtzia blanda</i> Benth. (Labiatae). <i>J. Ess. Oils Res.</i> 4: 121-124
<i>Rosmarinus x mendizabalii</i>	Shoot	0.1	33.0	-0.2591299820046205	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the <i>Rosmarinus eriocalyx-tomentosus</i> Complex in Southeast Spain. <i>J. Ess. Oil Res.</i> 5: 243-246.
<i>Rosmarinus x lavandulaceus</i>	Shoot	0.1	30.0	-0.29725541836979963	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the <i>Rosmarinus eriocalyx-tomentosus</i> Complex in Southeast Spain. <i>J. Ess. Oil Res.</i> 5: 243-246.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Origanum vulgare	Plant		25.0	-0.4702361796243996	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum vulgare	Plant		25.0	-0.4702361796243996	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Rosmarinus officinalis	Shoot	12.0	25.0	-0.3607978123117648	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Sassafras albidum	Leaf	9.4	21.0	0.24797623068876656	--
Rosmarinus tomentosus	Shoot	0.1	20.0	-0.4243402062537299	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Micromeria fruticosa	Shoot		19.0	-0.43704868504212296	Fleisher, Z. and Fleisher, A. 1991. The Essential Oil of Micromeria fruticosa (L.) Druce subsp. barbata (Boiss et. Ky.), P.H. Davis. Aromatic Plants of the Holy Land and the Sinai. Part VII. J. Ess. Oil Res 3: 477-479.
Micromeria fruticosa	Shoot		19.0	-0.43704868504212296	Fleisher, Z. and Fleisher, A. 1991. The Essential Oil of Micromeria fruticosa (L.) Druce subsp. barbata (Boiss et. Ky.), P.H. Davis. Aromatic Plants of the Holy Land and the Sinai. Part VII. J. Ess. Oil Res 3: 477-479.
Minthostachys mollis	Shoot	3.0	18.0	-0.449757163830516	Alkire, B.H., Tucker, A.O., and Maciarello, M.J. 1994. Tipo (Minthostachys mollis (Lamiaceae): An Ecuadorian Mint. Econ. Bot. 48(1): 60-64.
Origanum sipyleum	Shoot		17.0	-0.46246564261890905	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Rosmarinus officinalis	Shoot	0.1	13.0	-0.5132995577724812	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot		13.0	-0.5132995577724812	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Apium graveolens	Seed	7.0	10.0		Charalambous, G. (Ed.). 1994. Spices, Herbs and Edible Fungi. Elsevier Science B. V. Amsterdam. 764 pp.
Eucalyptus bridgesiana	Leaf		8.0	-0.6116747023656242	Singh, A. K., Gupta, K. C., & Brophy, J. J. 1991. Volatile Constituents of the Essential Oil of Eucalyptus bridgesiana Growing in India. Journal of Essential Oil Res. 3: 449-450.
Citrus paradisi	Pericarp		8.0		--
Dictamnus albus	Shoot		7.0	-0.5895504305028394	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Planta Med. 60:481-2
Dictamnus albus	Shoot		6.0	-0.6022589092912324	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Planta Med. 60:481-2
Dictamnus albus	Shoot		6.0	-0.6022589092912324	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Planta Med. 60:481-2
Citrus limon	Petiole		6.0		--
Origanum minutiflorum	Shoot	0.0	5.0	-0.6149673880796254	Baser, K.H.C., Tumen, G., Sezik, E. 1991. The Essential Oil of Origanum minutiflorum O. Schwarz and P.H. Davis. J. Ess. Oil Res. 3: 445-446.
Citrus reticulata	Fruit	4.0	5.0	-0.6151691947960529	--
Porophyllum ruderale	Plant	3.0	4.0	-0.608347505108489	Loayza, I., de Groot, W., Lorenzo, D. et al. 1999. Composition of the essential oil of Porophyllum ruderale (Jacq.) Cass. from Bolivia. Flav. & Fragr. J. 14: 393-8.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Leonotis leonurus	Se		3.0		Pedro, L.G., Barroso, J.G., Marques, N.T., Ascensao, L., Pais, M.S.S. and Scheffer, J.J.C. 1991. Composition of the Essential Oil from Sepals of Leonotis leonurus R. Br. J. Ess. Oil Res. 3: 451-3
Origanum sipyleum	Shoot		1.8	-0.6556345202024831	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Citrus sinensis	Fruit	0.7	1.0	-0.795218715224166	--
Pelargonium citrosum	Shoot		1.0	-0.6658013032331975	Matsuda, B. M., et al. 1996. Essential Oil Analysis and Field Evaluation of the Citrosa Plant 'Pelargonium citrosum' as a Repellent Against Populations of Aedes Mosquitoes. J. Am. Mosq. Contr. Assoc. 12(1):69-74.
Micromeria myrtifolia	Shoot		0.3	-0.6746972383850727	Ozek, T., Kirimer, N., and Baser, K.H.C. 1992. Composition of the Essential Oil of Micromeria myrtifolia Boiss. et Hohen. J. Ess. Oil Res., 4: 79-80.
Cinnamomum verum	Leaf		0.0	-1.1406906611683263	Mallavarupu, G. R. et al. 1995. Investigation of the essential oil of cinnamon leaf grown at Bangalore and Hyderabad. Flav. & Fragr. J., 10: 239-242.
Hyptis suaveolens	Shoot		0.0	-0.6785097820215906	Mallvarapu, G.R., Ramesh, S., Kaul, P.N., Bhattacharya, A.K., and Rao, B.R.R. 1993. The Essential Oil of Hyptis suaveolens (L.) Poit. J. Ess. Oil Res. 5: 321.
Hyptis suaveolens	Shoot		0.0	-0.6785097820215906	Mallvarapu, G.R., Ramesh, S., Kaul, P.N., Bhattacharya, A.K., and Rao, B.R.R. 1993. The Essential Oil of Hyptis suaveolens (L.) Poit. J. Ess. Oil Res. 5: 321.
Carum carvi	Seed				--
Carica papaya	Fruit				--