

List of Plants



Chemid

VIRIDIFLOROL

*Unless otherwise noted all references are to Duke, James A. 1992. Handbook of phytochemical constituents of GRAS herbs and other economic plants. Boca Raton, FL. CRC Press.

Plant	Part	Low PPM	High PPM	StdDev	*Reference
Acinos alpinus	Shoot	--	14.0	-0.79	Velasco-Negueruela,A., Perez-Alonso,M.J., Jiminez,S.M. and Garcia,F.M. 1993. The Volatile Constituents of Acinos alpinus (L.) Moench ssp. meridionalis (Nyman). P.W. Ball Growing in Spain. Flav. & Frag. J. 8:127-130.)
Acinos alpinus	Shoot	--	14.0	-0.79	Velasco-Negueruela,A., Perez-Alonso,M.J., Jiminez,S.M. and Garcia,F.M. 1993. The Volatile Constituents of Acinos alpinus (L.) Moench ssp. meridionalis (Nyman). P.W. Ball Growing in Spain. Flav. & Frag. J. 8:127-130.)
Cistus ladaniferus	Resin, Exudate, Sap	--	--		*
Eucalyptus melanophloia	Leaf	--	110.0	-0.21	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus Symphyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364
Eucalyptus behriana	Leaf	--	60.0	-0.44	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus Symphyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364
Eucalyptus leucoxylo	Leaf	--	175.0	0.09	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus Symphyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364
Eucalyptus globulus	Leaf Essent. Oil	--	--		*
Eucalyptus tetraptera	Leaf	--	210.0	0.25	Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus.Part I.Subgenus Symphyomyrtus,Section Dumaria,Series Incrassatae.Flavour and Fragrance J.9(3):113-7
Eucalyptus desquamata	Leaf	--	85.0	-0.33	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus Symphyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364
Eucalyptus intertexta	Leaf	--	14.0	-0.65	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus Symphyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364
Eucalyptus forrestiana	Leaf	--	210.0	0.25	Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus.Part I.Subgenus Symphyomyrtus,Section Dumaria,Series Incrassatae.Flavour and Fragrance J.9(3):113-7
Eucalyptus sparsa	Leaf	--	125.0	-0.14	*
Eucalyptus ceratocorys	Leaf	--	26.0	-0.6	Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus.Part I.Subgenus Symphyomyrtus,Section Dumaria,Series Incrassatae.Flavour and Fragrance J.9(3):113-7
Eucalyptus nova-anglica	Leaf	25.0	335.0	0.83	Brophy, J. L., Lassak, E. V., & Boland, D. J. 1992. The Leaf Essential Oils of Eucalyptus nova-anglica. Deane & Maiden. Journal of Essential Oil Res. 4: 29-32.
Eucalyptus lansdowneana	Leaf	--	160.0	0.02	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus Symphyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364

<i>Eucalyptus fasciculosa</i>	Leaf	--	3.0	-0.71	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> . Part VI. Subgenus <i>Symphyomyrtus</i> , Section <i>Adnataria</i> . Flavour and Fragrance J. 10(6):359-364
<i>Eucalyptus stoatei</i>	Leaf	--	250.0	0.44	Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> .Part I.Subgenus <i>Symphyomyrtus</i> ,Section <i>Dumaria</i> ,Series <i>Incrassatae</i> .Flavour and Fragrance J.9(3):113-7
<i>Eucalyptus odorata</i>	Leaf	--	18.0	-0.64	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> . Part VI. Subgenus <i>Symphyomyrtus</i> , Section <i>Adnataria</i> . Flavour and Fragrance J. 10(6):359-364
<i>Eucalyptus cuprea</i>	Leaf	--	70.0	-0.4	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> . Part VI. Subgenus <i>Symphyomyrtus</i> , Section <i>Adnataria</i> . Flavour and Fragrance J. 10(6):359-364
<i>Eucalyptus erythrandra</i>	Leaf	--	75.0	-0.37	Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> .Part I.Subgenus <i>Symphyomyrtus</i> ,Section <i>Dumaria</i> ,Series <i>Incrassatae</i> .Flavour and Fragrance J.9(3):113-7
<i>Eucalyptus populnea</i>	Leaf	--	60.0	-0.44	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> . Part VI. Subgenus <i>Symphyomyrtus</i> , Section <i>Adnataria</i> . Flavour and Fragrance J. 10(6):359-364
<i>Eucalyptus bridgesiana</i>	Leaf	--	60.0	-0.44	Singh, A. K., Gupta, K. C., & Brophy, J. J. 1991. Volatile Constituents of the Essential Oil of <i>Eucalyptus bridgesiana</i> Growing in India. Journal of Essential Oil Res. 3: 449-450.
<i>Eucalyptus angulosa</i>	Leaf	--	15.0	-0.65	Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> .Part I.Subgenus <i>Symphyomyrtus</i> ,Section <i>Dumaria</i> ,Series <i>Incrassatae</i> .Flavour and Fragrance J.9(3):113-7
<i>Eucalyptus globulus</i>	Leaf	--	--		Leung, A.Y., Encyclopedia of Common Natural Ingredients Used in Food, Drugs, and Cosmetics, John Wiley & Sons, New York, 1980.
<i>Eucalyptus ochrophloia</i>	Leaf	--	6.0	-0.69	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> . Part VI. Subgenus <i>Symphyomyrtus</i> , Section <i>Adnataria</i> . Flavour and Fragrance J. 10(6):359-364
<i>Eucalyptus viridis</i>	Leaf	--	13.0	-0.66	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> . Part VI. Subgenus <i>Symphyomyrtus</i> , Section <i>Adnataria</i> . Flavour and Fragrance J. 10(6):359-364
<i>Eucalyptus porosa</i>	Leaf	--	540.0	1.78	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> . Part VI. Subgenus <i>Symphyomyrtus</i> , Section <i>Adnataria</i> . Flavour and Fragrance J. 10(6):359-364
<i>Eucalyptus incrassata</i>	Leaf	--	295.0	0.65	Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> .Part I.Subgenus <i>Symphyomyrtus</i> ,Section <i>Dumaria</i> ,Series <i>Incrassatae</i> .Flavour and Fragrance J.9(3):113-7
<i>Eucalyptus largisparsa</i>	Leaf	--	110.0	-0.21	Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> . Part VI. Subgenus <i>Symphyomyrtus</i> , Section <i>Adnataria</i> . Flavour and Fragrance J. 10(6):359-364
<i>Eucalyptus dolichorhyncha</i>	Leaf	--	110.0	-0.21	Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus <i>Eucalyptus</i> .Part I.Subgenus <i>Symphyomyrtus</i> ,Section <i>Dumaria</i> ,Series <i>Incrassatae</i> .Flavour and Fragrance J.9(3):113-7
<i>Eucalyptus brassiana</i>	Leaf	--	25.0	-0.6	Singh, A. K., Gupta, K. C., & Brophy, J. J. 1991. Chemical Constituents of the Leaf Essential Oil of <i>Eucalyptus brassiana</i> S. T. Blake. Journal of Essential Oil Res. 3: 45-7.
<i>Humulus lupulus</i>	Fruit	--	16.0		*
<i>Juniperus oxycedrus</i>	Wood	--	--		List, P.H. and Horhammer, L., Hager's Handbuch der Pharmazeutischen Praxis, Vols. 2-6, Springer-Verlag, Berlin, 1969-1979.
<i>Laurus nobilis</i>	Leaf	--	--		Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
<i>Lycopus uniflorus</i>	Plant	--	2.0		Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.

Melaleuca alternifolia	Essential Oil	--	5800.0		*
Melaleuca alternifolia	Leaf	--	--		*
Melaleuca leucadendra	Plant	--	--		*
Mentha spicata	Leaf	--	--		Duke, J. A. Writeups or information summaries on approximately 2,000 economic plants, USDA, ARS, Beltsville, MD 20705.
Mentha aquatica	Shoot	--	140.0	1.11	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogekagaku Kaishi 67(10): 1417-1419.
Mentha aquatica	Shoot	--	140.0	1.11	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogekagaku Kaishi 67(10): 1417-1419.
Mentha longifolia	Shoot	--	85.0	0.28	Planta Medica, 57: A83, 1991.
Mentha aquatica	Shoot	--	195.0	1.94	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogekagaku Kaishi 67(10): 1417-1419.
Mentha aquatica	Leaf	36.0	1105.0	4.4	*
Mentha x piperita	Leaf	1.0	15.0	-0.65	Duke, J. A. Writeups or information summaries on approximately 2,000 economic plants, USDA, ARS, Beltsville, MD 20705.
Mentha aquatica	Shoot	--	50.0	-0.25	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogekagaku Kaishi 67(10): 1417-1419.
Mentha aquatica	Shoot	--	135.0	1.03	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogekagaku Kaishi 67(10): 1417-1419.
Mentha spicata	Essential Oil	--	--		*
Metrosideros sclerocarpa	Leaf	56.0	224.0	0.32	*
Salvia officinalis	Essential Oil	--	--		*
Salvia officinalis	Leaf	--	--		*
Tagetes minuta	Flower Essent. Oil	--	500.0		*
Tagetes lucida	Shoot	--	--		Bicchi, C., Fresia, M., Rubiolo, P., Monti, D., Franz, C., Goehler, I. 1997. Constituents of Tagetes lucida Cav. ssp. lucida essential oil. Flavor & Fragrance, 12(1): 47-52.
Teucrium polium	Shoot	--	3.5	-0.95	Perez-Alonso, M.J. Velasco-Negueruela, A. and Lopez-Saez, J.A. 1993. The Essential Oils of Two Iberian Teucrium Species. J. Ess. Oil Res. 5: 397-402.
Teucrium gnaphalodes	Shoot	--	4.0	-0.94	Perez-Alonso, M.J. Velasco-Negueruela, A. and Lopez-Saez, J.A. 1993. The Essential Oils of Two Iberian Teucrium Species. J. Ess. Oil Res. 5: 397-402.
Thymus funkii	Shoot	--	8.0	-0.88	Vila, R., et al. 1995. Composition and study of the variability of the essential oil of Thymus funkii Cousson. Flav. & Fragr. J. 10(6): 379-383.
Thymus funkii	Shoot	--	8.0	-0.88	Vila, R., et al. 1995. Composition and study of the variability of the essential oil of Thymus funkii Cousson. Flav. & Fragr. J. 10(6): 379-383.