

List of Plants

(Z)-BETA-OCIMENE

Chemid

ZBETAOCIMENE

*Unless otherwise noted all references are to Duke, James A. 1992. Handbook of phytochemical constituents of GRAS herbs and other economic plants. Boca Raton, FL. CRC Press.

Plant	Part	Low PPM	High PPM	StdDev	*Reference
Agastache foeniculum	Plant	3.0	120.0	1.41	*
Alpinia galanga	Rhizome	8.0	985.0		*
Alpinia galanga	Leaf	--	1705.0	1.41	*
Apium graveolens	Seed	6.0	9.0		Charalambous, G. (Ed.). 1994. Spices, Herbs and Edible Fungi. Elsevier Science B. V. Amsterdam. 764 pp.
Calamintha nepeta	Shoot	--	9.0	-0.41	Kirimer, N., Baser, K.H.C., Ozek, T. and Kurkcuoglu, M. 1992. Composition of the Essential Oil of Calamintha nepeta subsp. glandulosa. J. Ess. Oil Res. 4:189-190
Carica papaya	Fruit	--	--		*
Carum carvi	Seed	--	--		*
Cinnamomum verum	Leaf	--	--	-0.71	Mallavarupu, G. R. et al. 1995. Investigation of the essential oil of cinnamon leaf grown at Bangalore and Hyderabad. Flav. & Fragr. J., 10: 239-242.
Citrus paradisi	Pericarp	--	1.0		*
Citrus sinensis	Fruit	2.0	3.0	-0.66	*
Citrus limon	Petiole	--	3.5		*
Citrus reticulata	Fruit	--	0.1	-0.75	*
Dictamnus albus	Shoot	--	15.0	-0.38	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Planta Med. 60:481-2
Dictamnus albus	Shoot	--	15.0	-0.38	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Planta Med. 60:481-2
Dictamnus albus	Shoot	--	0.5	-0.45	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Planta Med. 60:481-2
Elsholtzia blanda	Shoot	--	18.0	-0.36	Bestman, H.J., Rauscher, J., Vostrowsky O., Pant, A.K., Dev. V., Perihar, R. and Mathela, C.S. 1992. Constituents of the Essential Oil of Elsholtzia blanda Benth. (Labiatae). J. Ess. Oils Res. 4: 121-124
Hyptis suaveolens	Shoot	--	3.0	-0.43	Mallvarapu, G.R., Ramesh, S., Kaul, P.N., Bhattacharya, A.K., and Rao, B.R.R. 1993. The Essential Oil of Hyptis suaveolens (L.) Poit. J. Ess. Oil Res. 5: 321.
Hyptis suaveolens	Shoot	--	3.0	-0.43	Mallvarapu, G.R., Ramesh, S., Kaul, P.N., Bhattacharya, A.K., and Rao, B.R.R. 1993. The Essential Oil of Hyptis suaveolens (L.) Poit. J. Ess. Oil Res. 5: 321.
Hyssopus officinalis	Shoot	--	980.0	4.06	Tsankova, E.T., Konatchiev, A.N. and Genova, E.M. 1993. Chemical Composition of the Essential Oils of Two Hyssopus officinalis cultivars. J. Ess. Oil Res. 5: 609-611.
Hyssopus officinalis	Shoot	--	690.0	2.72	Tsankova, E.T., Konatchiev, A.N. and Genova, E.M. 1993. Chemical Composition of the Essential Oils of Two Hyssopus officinalis cultivars. J. Ess. Oil Res. 5: 609-611.
Leonotis leonurus	Se	--	24.0		Pedro, L.G., Barroso, J.G., Marques, N.T., Ascensao, L., Pais, M.S.S. and Scheffer, J.J.C. 1991. Composition of the Essential Oil from Sepals of Leonotis leonurus R. Br. J. Ess. Oil Res. 3: 451-3
Lindera benzoin	Fruit	55.0	75.0	1.41	*
Minthostachys mollis	Shoot	4.0	8.0	-0.41	Alkire, B.H., Tucker, A.O., and Maciarelo, M.J. 1994. Tipo (Minthostachys mollis (Lamiaceae): An Ecuadorian Mint. Econ. Bot. 48(1): 60-64.

Nepeta racemosa	Shoot	--	33.0	-0.3	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.
Nepeta racemosa	Shoot	--	33.0	-0.3	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.
Origanum vulgare	Plant	--	5.0	-0.78	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum vulgare	Plant	--	20.0	-0.5	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum sipyleum	Shoot	--	15.0	-0.38	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum vulgare	Plant	--	5.0	-0.78	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum sipyleum	Shoot	--	7.0	-0.42	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum sipyleum	Shoot	--	205.0	0.49	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum vulgare	Plant	--	7.0	-0.75	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum sipyleum	Shoot	--	205.0	0.49	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Pelargonium citrosum	Shoot	--	3.0	-0.43	Matsuda, B. M., et al. 1996. Essential Oil Analysis and Field Evaluation of the Citrosa Plant 'Pelargonium citrosum' as a Repellent Against Populations of Aedes Mosquitoes. J. Am. Mosq. Contr. Assoc. 12(1):69-74.
Porophyllum ruderale	Plant	80.0	120.0	1.41	Loayza, I., de Groot, W., Lorenzo, D. et al. 1999. Composition of the essential oil of Porophyllum ruderale (Jacq.) Cass. from Bolivia. Flav. & Fragr. J. 14: 393-8.
Rosmarinus x mendizabali	Shoot	--	0.1	-0.45	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot	6.0	75.0	-0.1	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot	--	13.0	-0.39	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus x lavandulaceus	Shoot	26.0	110.0	0.06	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus eriocalyx	Shoot	--	--		Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus tomentosus	Shoot	--	0.1	-0.45	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot	--	--		Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus eriocalyx	Shoot	--	0.1	-0.45	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot	6.0	75.0	-0.1	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Sassafras albidum	Leaf	2.6	8.0	-0.7	*
Satureja cilicica	Shoot	--	210.0	0.52	Tumen, G. Baser, K.H.C. and Kirimer, N. 1993. The Essential Oil of Satureja cilicica P.H. Davis. J. Ess. Oil Res. 5: 547-548.

Tagetes lucida	Shoot	--	--		Bicchi, C., Fresia, M., Rubiolo, P., Monti, D., Franz, C., Goehler, I. 1997. Constituents of Tagetes lucida Cav. ssp. lucida essential oil. Flavor & Fragrance, 12(1): 47-52.
Thymus longicaulis	Shoot	--	0.8	-0.44	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot	--	0.8	-0.44	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot	--	--	-0.45	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.