

List of Plants

5-METHYL-3-HEPTANONE

Chemid

5METHYL3HEPTANONE

*Unless otherwise noted all references are to Duke, James A. 1992. Handbook of phytochemical constituents of GRAS herbs and other economic plants. Boca Raton, FL. CRC Press.

Plant	Part	Low PPM	High PPM	StdDev	*Reference
Nepeta racemosa	Shoot	--	52.0	1.58	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.
Nepeta racemosa	Shoot	--	52.0	1.58	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.
Sideritis athoa	Shoot	--	0.5	-0.77	Ozek, T., Baser, K.H.C. and Tumen, G. 1993. The Essential Oil of Sideritis athoa Papanikolaou Et Kokkini. J. Ess. Oil Res. 5: 669-670.
Thymus longicaulis	Shoot	--	5.0	-0.56	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot	--	3.0	-0.65	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot	--	5.0	-0.56	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus cilicicus	Shoot	--	4.0	-0.61	Tumen, G., Koyuncu, M., Kirimer, N., and Baser, K.H.C. 1994. Composition of the Essential Oil of Thymus cilicicus Boiss. & Bal. J. Ess. Oil Res. 6: 97-8.