

# List of Plants



## Chemid

BAICALIN

## Dosage

LD50=3,081 mg/kg (ivn mus) Leung, A. Y. and Foster, S. 1995. Encyclopedia of Common Natural Ingredients 2nd Ed. John Wiley & Sons, New York. 649 pp.

LDI0=15 mg/kg (ivn rbt) Leung, A. Y. and Foster, S. 1995. Encyclopedia of Common Natural Ingredients 2nd Ed. John Wiley & Sons, New York. 649 pp.

LD50=3,000 (ipr mus) Merck 11th Edition

\*Unless otherwise noted all references are to Duke, James A. 1992. Handbook of phytochemical constituents of GRAS herbs and other economic plants. Boca Raton, FL. CRC Press.

Plant	Part	Low PPM	High PPM	StdDev	*Reference
Plantago major	Plant	--	--		*
Scutellaria baicalensis	Tissue Culture	--	--		*
Scutellaria sp	Plant	--	--		Jeffery B. Harborne and H. Baxter, eds. 1983. Phytochemical Dictionary. A Handbook of Bioactive Compounds from Plants. Taylor & Frost, London. 791 pp.
Scutellaria baicalensis	Plant	--	--		*
Scutellaria galericulata	Leaf	--	10000.0		List, P.H. and Horhammer, L., Hager's Handbuch der Pharmazeutischen Praxis, Vols. 2-6, Springer-Verlag, Berlin, 1969-1979.
Scutellaria baicalensis	Stem	5800.0	43000.0		*
Scutellaria baicalensis	Root	1000.0	206000.0		*