

Dr. Duke's Phytochemical and Ethnobotanical Database

List of Plants for ALPHA-EUDESMOL

| Plant | Part | Low PPM | High PPM | StdDev | Reference |
|---------------------------|------|------------|-------------|----------------------|--|
| Apium graveolens | Seed | 76 | 225 | | -- |
| Eucalyptus angulosa | Leaf | -- | 100 | 0.023093885596796895 | Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus.Part I.Subgenus Sympyomyrtus,Section Dumaria,Series Incrassatae.Flavour and Fragrance J.9(3):113-7 |
| Eucalyptus behriana | Leaf | 15 | 15 | -0.6020590684566249 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus Sympyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Eucalyptus brassiana | Leaf | -- | 0 | -0.712380177995464 | Singh, A. K., Gupta, K. C., & Brophy, J. J. 1991. Chemical Constituents of the Leaf Essential Oil of Eucalyptus brassiana S. T. Blake. Journal of Essential Oil Res. 3: 45-7. |
| Eucalyptus ceratocorys | Leaf | -- | 50 | -0.34464314619933356 | Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus.Part I.Subgenus Sympyomyrtus,Section Dumaria,Series Incrassatae.Flavour and Fragrance J.9(3):113-7 |
| Eucalyptus cuprea | Leaf | 315 | 315 | 1.604363122320158 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus Sympyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Eucalyptus desquamata | Leaf | 22 | 22 | -0.5505758840051667 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus Sympyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Eucalyptus dolichorhyncha | Leaf | -- | 125 | 0.20696240149486214 | Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus.Part I.Subgenus Sympyomyrtus,Section Dumaria,Series Incrassatae.Flavour and Fragrance J.9(3):113-7 |

| Plant | Part | Low PPM | High PPM | StdDev | Reference |
|-------------------------|------------------------|---------|----------|---------------------|---|
| Eucalyptus erythrandra | Leaf | -- | 205 | 0.7953416523686709 | Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus.Part I.Subgenus Sympyomyrtus,Section Dumaria, Series Incrassatae.Flavour and Fragrance J.9(3):113-7 |
| Eucalyptus fasiculosa | Leaf | 17 | 17 | -0.5873495871847797 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subjenus Sympyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Eucalyptus forrestiana | Leaf | -- | 320 | 1.641136825499771 | Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus.Part I.Subgenus Sympyomyrtus,Section Dumaria, Series Incrassatae.Flavour and Fragrance J.9(3):113-7 |
| Eucalyptus globulus | Plant | -- | -- | | J.S. Glasby Dict.Pls Containing 2ndary Metabolite. 1991. |
| Eucalyptus globulus | Leaf Essent. Oil | -- | 17000 | | -- |
| Eucalyptus incrassata | Leaf | -- | 295 | 1.4572683096017058 | Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus.Part I.Subgenus Sympyomyrtus,Section Dumaria, Series Incrassatae.Flavour and Fragrance J.9(3):113-7 |
| Eucalyptus intertexta | Leaf | 5.5 | 5.5 | -0.6719291044978897 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subjenus Sympyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Eucalyptus lansdowneana | Leaf | 14 | 14 | -0.6094138090925475 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subjenus Sympyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Eucalyptus largisparsa | Leaf | 15 | 15 | -0.6020590684566249 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subjenus Sympyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |

| Plant | Part | Low PPM | High PPM | StdDev | Reference |
|-------------------------|-------|---------|----------|----------------------|---|
| Eucalyptus leucoxylon | Leaf | 270 | 270 | 1.2733997937036405 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus <i>Syphomyrtus</i> , Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Eucalyptus melanophloia | Leaf | 6 | 6 | -0.6682517341799284 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus <i>Syphomyrtus</i> , Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Eucalyptus nova-anglica | Leaf | 20 | 35 | -0.45496425573817273 | Brophy, J. L., Lassak, E. V., & Boland, D. J. 1992. The Leaf Essential Oils of Eucalyptus nova-anglica. Deane & Maiden. Journal of Essential Oil Res. 4: 29-32. |
| Eucalyptus ochrophloia | Leaf | 7 | 7 | -0.6608969935440058 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus <i>Syphomyrtus</i> , Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Eucalyptus odorata | Leaf | 25 | 25 | -0.5285116620973989 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus <i>Syphomyrtus</i> , Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Eucalyptus populnea | Leaf | 3 | 3 | -0.6903159560876962 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus <i>Syphomyrtus</i> , Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Eucalyptus porosa | Leaf | 13 | 13 | -0.6167685497284702 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus <i>Syphomyrtus</i> , Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Eucalyptus sp | Plant | -- | -- | | Jeffery B. Harborne and H. Baxter, eds. 1983. Phytochemical Dictionary. A Handbook of Bioactive Compounds from Plants. Taylor & Frost, London. 791 pp. |
| Eucalyptus sparsa | Leaf | 25 | 25 | -0.5285116620973989 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subgenus <i>Syphomyrtus</i> , Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |

| Plant | Part | Low PPM | High PPM | StdDev | Reference |
|-----------------------|---------------|---------|----------|---------------------|---|
| Eucalyptus stoatei | Leaf | -- | 510 | 3.0385375463250672 | Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus.Part I.Subgenus Sympyomyrtus,Section Dumaria, Series Incrassatae.Flavour and Fragrance J.9(3):113-7 |
| Eucalyptus tetraptera | Leaf | -- | 27 | -0.5138021808255536 | Bignell,C.M.,Dunlop,P.J.,Brophy,J.J.,and Jackson, J.F.1994.Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus.Part I.Subgenus Sympyomyrtus,Section Dumaria, Series Incrassatae.Flavour and Fragrance J.9(3):113-7 |
| Eucalyptus viridis | Leaf | 2 | 2 | -0.6976706967236188 | Bignell, C.M., Dunlop, P.J., Brophy, J.J., and Jackson, J.F. 1995. Volatile Leaf Oils of Some South-western and Southern Australian Species of the Genus Eucalyptus. Part VI. Subjenus Sympyomyrtus, Section Adnataria. Flavour and Fragrance J. 10(6):359-364 |
| Ferula gummosa | Plant | -- | -- | | -- |
| Humulus lupulus | Fruit | 65 | 65 | | -- |
| Hypericum perforatum | Plant | 2.5 | 45 | | -- |
| Juniperus virginiana | Leaf | -- | -- | | -- |
| Magnolia officinalis | Essential Oil | -- | -- | | -- |
| Mentha aquatica | Shoot | 7 | 7 | -0.3517412136650936 | Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419. |
| Mentha aquatica | Shoot | -- | 22 | 1.3270236697364897 | Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419. |
| Mentha aquatica | Shoot | -- | 24 | 1.5508589875233674 | Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419. |

| Plant | Part | Low PPM | High PPM | StdDev | Reference |
|-----------------------|-------|---------|----------|---------------------|--|
| Mentha aquatica | Shoot | -- | 11 | 0.09592942190866195 | Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419. |
| Mentha aquatica | Shoot | -- | 7 | -0.3517412136650936 | Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419. |
| Rhododendron dauricum | Plant | -- | -- | | -- |
| Thuja occidentalis | Wood | -- | -- | | -- |
| Thymus funkii | Shoot | 0 | 0 | -1.1351648259191658 | Vila, R., et al. 1995. Composition and study of the variability of the essential oil of <i>Thymus funkii</i> Cousson. <i>Flav. & Fragr. J.</i> 10(6): 379-383. |
| Thymus funkii | Shoot | -- | 0 | -1.1351648259191658 | Vila, R., et al. 1995. Composition and study of the variability of the essential oil of <i>Thymus funkii</i> Cousson. <i>Flav. & Fragr. J.</i> 10(6): 379-383. |