

# C BOURBONENE

## Chemid

BOURBONENE

\*Unless otherwise noted all references are to Duke, James A. 1992. Handbook of phytochemical constituents of GRAS herbs and other economic plants. Boca Raton, FL. CRC Press.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Cymbopogon nardus	Plant	30	120		Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
Cymbopogon winterianus	Plant	--	--		Guenther, E., The Essential Oils, 6 volumes, D. van Nostrand, New York, 1948-1952.
Elsholtzia cristata	Shoot	11	11	2.0308775244492967	Kobold, U., Vostrowsky, O., Bestmann, H.J., Bisht, J.C., Pant, A.K., Melkani, A.B. and Mathela, C.S. 1987. Terpenoids from Elsholtzia Species; II. Constituents of Essential Oil from a New Chemotype of Elsholtzia cristata. Planta Medica 1987: 268-271.
Mentha aquatica	Shoot	6	6	0.0655121782080416	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
Mentha aquatica	Shoot	--	4	-0.7206339602884604	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
Mentha aquatica	Shoot	--	3	-1.1137070295367113	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Mentha aquatica	Shoot --	5	-0.3275608910402094		Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
Mentha aquatica	Shoot --	6	0.0655121782080416		Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.