

C (E)-2-HEXENAL

Chemid

E2HEXENAL

CAS Number

6728263

*Unless otherwise noted all references are to Duke, James A. 1992. Handbook of phytochemical constituents of GRAS herbs and other economic plants. Boca Raton, FL. CRC Press.

Plant	Part	Low PPM	High PPM	StdDev PPM	Reference
Callicarpa americana	Leaf	--	50	0.9999999999999997	--
Carthamus tinctorius	Flower	0.5	0.6	-1.0000000000000002	--
Carthamus tinctorius	Leaf	1.67	1.67	-1	--
Diospyros virginiana	Fruit	--	--	--	--
Hibiscus sabdariffa	Calyx	--	--	--	--
Nepeta racemosa	Shoot	3	3	0.25537695922762477	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.

Plant	Part	Low PPM	High PPM	StdDev PPM	Reference
Origanum vulgare	Plant	0.5	0.5	-0.6708631792558644	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four <i>Origanum vulgare</i> Subspecies of Anatolian Origin. <i>J. Ess. Oil Res.</i> , 5: 425-431.
Origanum vulgare	Plant	15	15	1.4136045562891424	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four <i>Origanum vulgare</i> Subspecies of Anatolian Origin. <i>J. Ess. Oil Res.</i> , 5: 425-431.
Origanum vulgare	Plant	0	0	-0.7427413770332784	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four <i>Origanum vulgare</i> Subspecies of Anatolian Origin. <i>J. Ess. Oil Res.</i> , 5: 425-431.
Satureja cilicica	Shoot	5	5	1.5322617553657476	Tumen, G. Baser, K.H.C. and Kirimer, N. 1993. The Essential Oil of <i>Satureja cilicica</i> P.H. Davis. <i>J. Ess. Oil Res.</i> 5: 547-548.
Sideritis athoa	Shoot	4	4	0.8938193572966862	Ozek, T., Baser, K.H.C. and Tumen, G. 1993. The Essential Oil of <i>Sideritis athoa</i> Papanikolaou Et Kokkini. <i>J. Ess. Oil Res.</i> 5: 669-670.
Thymus longicaulis	Shoot	0.8	0.8	-1.1491963165243104	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of <i>Thymus longicaulis</i> C. Presl subsp. <i>longicaulis</i> in the same Population. <i>J. Ess. Oil Res.</i> 5: 291-5.
Thymus longicaulis	Shoot	--	2	-0.3830654388414366	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of <i>Thymus longicaulis</i> C. Presl subsp. <i>longicaulis</i> in the same Population. <i>J. Ess. Oil Res.</i> 5: 291-5.
Thymus longicaulis	Shoot	--	0.8	-1.1491963165243104	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of <i>Thymus longicaulis</i> C. Presl subsp. <i>longicaulis</i> in the same Population. <i>J. Ess. Oil Res.</i> 5: 291-5.
Trifolium pratense	Flower	--	20	1	Buchnauer,G.,Jirovetz,L.,Nikiforov,A.1996.Comparative Investigation of Essential Clover Flower Oils from Austria Using Gas Chromatography-Flame Ionization Detection, Gas Chromatography-Mass Spectrometry, and Gas Chrom.-Olfactometry. <i>J.Agrc.Food Chem.</i> 44:1827-8