

C T-CADINOL

Chemid

TCADINOL

*Unless otherwise noted all references are to Duke, James A. 1992. Handbook of phytochemical constituents of GRAS herbs and other economic plants. Boca Raton, FL. CRC Press.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Achillea millefolium	Leaf	1	15	-0.5225815390117998	--
Acinos alpinus	Shoot	2.5	2.5	-0.8937792727642405	Velasco-Negueruela,A., Perez-Alonso,M.J., Jiminez,S.M. and Garcia,F.M. 1993. The Volatile Constituents of Acinus alpinus (L.) Moench ssp. meridionalis (Nyman). P.W. Ball Growing in Spain. Flav. & Frag. J. 8:127-130.)
Acinos alpinus	Shoot	--	2.5	-0.8937792727642405	Velasco-Negueruela,A., Perez-Alonso,M.J., Jiminez,S.M. and Garcia,F.M. 1993. The Volatile Constituents of Acinus alpinus (L.) Moench ssp. meridionalis (Nyman). P.W. Ball Growing in Spain. Flav. & Frag. J. 8:127-130.)
Aloysia citrodora	Plant	5	35	-0.23187492634055548	Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
Artemisia pallens	Plant	38	168	1.6955853988653122	--
Cinnamomum camphora	Plant	--	--		--
Humulus lupulus	Fruit	9	9		--
Hyssopus officinalis	Leaf	0.1	60	0.2141790245785401	Flavour and Fragrance Journal, 6: 72.
Hyssopus officinalis	Shoot	30	30	0.6871033072491771	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Hyssopus officinalis	Shoot	--	20	0.1122369145170252	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot	--	30	0.6871033072491771	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot	--	30	0.6871033072491771	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot	--	--		Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot	--	--		Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot	--	30	0.6871033072491771	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot	--	--		Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot	--	20	0.1122369145170252	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Hyssopus officinalis	Shoot	--	30	0.6871033072491771	Kerrola, K., Galambosi, B. and Kallio, H. 1994. Volatile Components and Odor Intensity of Four Phenotypes of Hyssop (Hyssopus officinalis L.) J. Agric. Food Chem. 42: 776-781.
Melia azedarach	Wood	--	--		--

Plant	Part	Low PPM	High PPM	StdDev	Reference
Melissa officinalis	Shoot	3	24	0.342183471609886	Deutsche Apot. Ztit. 129(4):155-163. W. Schulze et al. Die Melisse.
Metrosideros nervulosa	Leaf	176	176	2.1133840329447495	--
Nepeta racemosa	Shoot	60	60	2.411702485445633	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.
Nepeta racemosa	Shoot	--	60	2.411702485445633	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.
Ocimum basilicum	Plant	0.5	0.5	-0.7318552362623783	Die Nahrung. Pino, J., Rosado, A., Goire, I., Roncal, E., and Garcia, I. 1993. Analysis of the Essential Oil from Cuban Basil. Die Nahrung 37:(5): 501-504.
Ocimum basilicum	Plant	--	0.5	-0.7318552362623783	Die Nahrung. Pino, J., Rosado, A., Goire, I., Roncal, E., and Garcia, I. 1993. Analysis of the Essential Oil from Cuban Basil. Die Nahrung 37:(5): 501-504.
Ocimum basilicum	Shoot Essent. Oil	--	71400		--
Pinus sylvestris	Leaf Essent. Oil	--	2000	-1	--
Pinus sylvestris	Leaf	--	--		Leung, A. Y. and Foster, S. 1995. Encyclopedia of Common Natural Ingredients 2nd Ed. John Wiley & Sons, New York. 649 pp.
Plectranthus barbatus	Essential Oil	5000	41000		--
Salvia dorisiana	Shoot	1.2	1.8	-0.9340199202554911	Tucker, A.O. & Maciarelo, M.J. 1994. The Essential Oil of Salvia dorisiana Standley. J. Ess. Oil Res. 6: 97-8.
Satureja obovata	Leaf	0.01	0.01	-0.7680046689722265	Arrebola, M.L., Navaro, M.C., Jimenez, J. and Ocana, F.A. 1994. Variations in Yield and Composition of the Essential Oil of Satureja obovata. Phytochemistry 35(1): 83.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Satureja obovata	Leaf	--	--		Arrebola, M.L., Navaro, M.C., Jimenez, J. and Ocana, F.A. 1994. Variations in Yield and Composition of the Essential Oil of Satureja obovata. <i>Phytochemistry</i> 35(1): 83.
Satureja obovata	Leaf	--	--		Arrebola, M.L., Navaro, M.C., Jimenez, J. and Ocana, F.A. 1994. Variations in Yield and Composition of the Essential Oil of Satureja obovata. <i>Phytochemistry</i> 35(1): 83.
Satureja obovata	Leaf	--	--		Arrebola, M.L., Navaro, M.C., Jimenez, J. and Ocana, F.A. 1994. Variations in Yield and Composition of the Essential Oil of Satureja obovata. <i>Phytochemistry</i> 35(1): 83.
Stevia rebaudiana	Flower	--	60		Kinghorn, A. D. (Ed.) 2002. Medicinal and Aromatic Plants - Industrial Profiles. Stevia. The genus Stevia. Taylor & Francis. New York, NY. 211 pp.
Stevia rebaudiana	Leaf	--	28	-0.309739598419035	Kinghorn, A. D. (Ed.) 2002. Medicinal and Aromatic Plants - Industrial Profiles. Stevia. The genus Stevia. Taylor & Francis. New York, NY. 211 pp.
Tagetes lucida	Shoot	--	6	-0.6925760353079873	Bicchi, C., Fresia, M., Rubiolo, P., Monti, D., Franz, C., Goehler, I. 1997. Constituents of Tagetes lucida Cav. ssp. lucida essential oil. <i>Flavor & Fragrance</i> , 12(1): 47-52.
Teucrium asiaticum	Shoot	1.16	1.16	-0.9708113693903488	Velasco-Negueruela, A. and Perez-Alonso, M.J. 1990. The Volatiles of Six Teucrium Species from the Iberian Peninsula and the Balearic Islands. <i>Phytochemistry</i> 29(4): 1165-9.
Teucrium gnaphalodes	Shoot	2.5	2.5	-0.8937792727642405	Perez-Alonso, M.J. Velasco-Negueruela, A. and Lopez-Saez, J.A. 1993. The Essential Oils of Two Iberian Teucrium Species. <i>J. Ess. Oil Res.</i> 5: 397-402.
Teucrium oxylepis	Shoot	9.24	9.24	-0.5063193240627701	Velasco-Negueruela, A. and Perez-Alonso, M.J. 1990. The Volatiles of Six Teucrium Species from the Iberian Peninsula and the Balearic Islands. <i>Phytochemistry</i> 29(4): 1165-9.
Teucrium oxylepis	Shoot	1	1	-0.9800092316740633	Velasco-Negueruela, A. and Perez-Alonso, M.J. 1990. The Volatiles of Six Teucrium Species from the Iberian Peninsula and the Balearic Islands. <i>Phytochemistry</i> 29(4): 1165-9.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Teucrium polium	Shoot	3	3	-0.8650359531276329	Perez-Alonso, M.J. Velasco-Negueruela, A. and Lopez-Saez, J.A. 1993. The Essential Oils of Two Iberian Teucrium Species. J. Ess. Oil Res. 5: 397-402.
Teucrium pseudoscorodonia	Shoot	0.82	0.82	-0.990356826743242	Velasco-Negueruela, A. and Perez-Alonso, M.J. 1990. The Volatiles of Six Teucrium Species from the Iberian Peninsula and the Balearic Islands. Phytochemistry 29(4): 1165-9.
Teucrium salviastrum	Shoot	0.65	0.65	-1.0001295554196887	Velasco-Negueruela, A. and Perez-Alonso, M.J. 1990. The Volatiles of Six Teucrium Species from the Iberian Peninsula and the Balearic Islands. Phytochemistry 29(4): 1165-9.
Teucrium scorodonia	Shoot	1.02	1.02	-0.978859498888599	Velasco-Negueruela, A. and Perez-Alonso, M.J. 1990. The Volatiles of Six Teucrium Species from the Iberian Peninsula and the Balearic Islands. Phytochemistry 29(4): 1165-9.
Thymus funkii	Shoot	35	35	0.9745365036152531	Vila, R., et al. 1995. Composition and study of the variability of the essential oil of Thymus funkii Cousson. Flav. & Fragr. J. 10(6): 379-383.
Thymus funkii	Shoot	--	25	0.39967011088310117	Vila, R., et al. 1995. Composition and study of the variability of the essential oil of Thymus funkii Cousson. Flav. & Fragr. J. 10(6): 379-383.
Thymus riatarum	Shoot	25	25	0.39967011088310117	Iglesias, J., Vila, R., Canigueral, S., Bellakdhar, and Il Idrissi, A. 1991. Analysis of the Essential Oil of Thymus riatarum. J. Ess. Oil Res. 3: 43-4.
Vitex agnus-castus	Leaf	--	2.5	-0.7272372511202277	Ekundayo, O., Laakso, I., Holopainen, M., Hiltunen, R., Oguntimein, B., and Kauppinen, V. 1990. The Chemical Composition and Antimicrobial Activity of the Leaf Oil of Vitex agnus-castus L. J. Essential Oil Research, 2: 115-119.
Vitex agnus-castus	Leaf Essent. Oil	--	10700	1	Jim Duke's personal files.
Vitex agnus-castus	Flower	--	--		Jim Duke's personal files.
Vitex agnus-castus	Fruit	--	--		Jim Duke's personal files.