

C 1-NONEN-3-OL

Chemid

1NONEN3OL

*Unless otherwise noted all references are to Duke, James A. 1992. Handbook of phytochemical constituents of GRAS herbs and other economic plants. Boca Raton, FL. CRC Press.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Origanum vulgare	Plant 1	1	1	1.3887301496588274	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum vulgare	Plant 0	0	0	-0.9258200997725514	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum vulgare	Plant 0.2	0.2	0.2	-0.46291004988627565	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Satureja cilicica	Shoot 6	6	6	-0.4642204280759514	Tumen, G. Baser, K.H.C. and Kirimer, N. 1993. The Essential Oil of Satureja cilicica P.H. Davis. J. Ess. Oil Res. 5: 547-548.
Sideritis scardica	Shoot 110	110	110	1.9989900066127695	Menkovic, N., et al. 1991. The Essential Oil of Sideritis scardica. Pl. Med. 57. Suppl. 2. pp. A137-A132.
Thymus longicaulis	Shoot 5	5	5	-0.48790514379411215	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot --	2	2	-0.5589592909485944	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot --	5	5	-0.48790514379411215	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.