

Dr. Duke's Phytochemical and Ethnobotanical Database

List of Plants for 6-METHYL-3-HEPTANOL

Plant	Part	Low PPM	High PPM	StdDev	Reference
Calamintha nepeta	Plant	75	75	2.2317566235144057	Kirimer, N., Baser, K.H.C., Ozek, T. and Kurkcuoglu, M. 1992. Composition of the Essential Oil of Calamintha nepeta subsp. glandulosa. J. Ess. Oil Res. 4:189-190
Micromeria myrtifolia	Shoot	0.1	0.1	-1.1458737449874261	Ozek, T., Kirimer, N., and Baser, K.H.C. 1992. Composition of the Essential Oil of Micromeria myrtifolia Boiss. et Hohen. J. Ess. Oil Res., 4: 79-80.
Nepeta racemosa	Shoot	13	13	0.5038775173270604	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.
Nepeta racemosa	Shoot	--	13	0.5038775173270604	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of Nepeta racemosa Lam. J. Ess. Oil Res. 5: 215-7.
Origanum vulgare	Plant	1	1	-0.5131185588689631	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum vulgare	Plant	1	1	-0.5131185588689631	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum vulgare	Plant	3	3	-0.43893274312887204	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum vulgare	Plant	3	3	-0.43893274312887204	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Satureja cilicica	Shoot	4	4	-0.6471117354504884	Tumen, G. Baser, K.H.C. and Kirimer, N. 1993. The Essential Oil of Satureja cilicica P.H. Davis. J. Ess. Oil Res. 5: 547-548.
Satureja cuneifolia	Shoot	4	4	-0.6471117354504884	Tumen, G. 1991. The Volatile Constituents of Satureja cuneifolia. J. Ess. Oil Res., 3: 365-366.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Sideritis athoa	Shoot	0.5	0.5	-1.094718667086202	Ozek, T., Baser, K.H.C. and Tumen, G. 1993. The Essential Oil of Sideritis athoa Papanikolaou Et Kokkini. J. Ess. Oil Res. 5: 669-670.
Sideritis germanicolpitana	Plant	5	6	-0.3276540195187355	J. Essential Oil, 4: 533.
Thymus cilicicus	Shoot	22	22	1.6548667701046094	Tumen, G., Koyuncu, M., Kirimer, N., and Baser, K.H.C. 1994. Composition of the Essential Oil of Thymus cilicicus Boiss. & Bal. J. Ess. Oil Res. 6: 97-8.
Thymus longicaulis	Shoot	17	17	1.0154282963393044	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot	--	0	-1.1586625144627323	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot	--	17	1.0154282963393044	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.