

# C (E)-NEROLIDOL

## Chemid

ENEROLIDOL

\*Unless otherwise noted all references are to Duke, James A. 1992. Handbook of phytochemical constituents of GRAS herbs and other economic plants. Boca Raton, FL. CRC Press.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Achillea millefolium	Leaf	--	--		--
Elsholtzia blanda	Shoot	2	2	-0.7716083190648096	Bestman, H.J., Rauscher, J., Vostrowsky O., Pant, A.K., Dev. V., Perihar, R. and Mathela, C.S. 1992. Constituents of the Essential Oil of Elsholtzia blanda Benth. (Labiatae). J. Ess. Oils Res. 4: 121-124
Lindera benzoin	Leaf	10	28		--
Lindera benzoin	Shoot Essent. Oil	--	100000		Tucker, A. O. and Debaggio, T. 2000. The Big Book of Herbs. Interweave Press, Inc. Loveland, CO. 688 pp.
Salvia dorisiana	Shoot	2.3	2.6	-0.6405804912990873	Tucker, A.O. & Maciarello, M.J. 1994. The Essential Oil of Salvia dorisiana Standley. J. Ess. Oil Res. 6: 97-8.
Tagetes lucida	Shoot	--	12	1.4121888103638975	Bicchi, C., Fresia, M., Rubiolo, P., Monti, D., Franz, C., Goehler, I. 1997. Constituents of Tagetes lucida Cav. ssp. lucida essential oil. Flavor & Fragrance, 12(1): 47-52.