

Dr. Duke's Phytochemical and Ethnobotanical Databases

Chemicals found in *Mentha aquatica*

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	ALPHA-EUDESOL	Shoot		7.0	-0.3517412136650936	Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	CIS-BETA-OCIMENE	Shoot		5.0	-0.5234943490109244	Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	1-OCTYL-3-YL-ACETATE	Shoot		3.0	-0.43804835651682184	Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-FARNESENE	Shoot		11.0	-0.3998562540982541	Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	MELISSIC-ACID	Leaf				List, P.H. and Horhammer, L., Hager's Handbuch der Pharmazeutischen Praxis, Vols. 2-6, Springer-Verlag, Berlin, 1969-1979.
0	N-HEXANAL	Shoot		2.0	1.0000000000000002	Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	HESPERETIN-7-BETA-D-GLUCOPYRANOSIDE	Plant				J.S. Glasby Dict.Pls Containing 2ndary Metabolite. 1991.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	VERBENYL-ACETATE	Shoot		2.0	-0.8164965809277258	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-CUBEBENE	Shoot		4.0	-0.253152042232558	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	GAMMA-MUROLENE	Shoot		1.0	-0.3724734981350398	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALLO-AROMADENDRENE	Shoot		3.0	-0.18318506682182947	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BOURBONENE	Shoot		5.0	-0.3275608910402094	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	EO	Shoot		1700.0	-0.7956845977843217	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	P-CYMOL	Leaf	6.0	51.0	-0.5242397055421542	--

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	2-N-AMYL FURAN	Shoot		0.1		Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALPHA-EUDES MOL	Shoot		11.0	0.09592942190866195	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	CIS-BETA-OCIMENE	Shoot		8.0	-0.5039521391496569	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	1-OCTEN-3-OL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-FARNESENE	Shoot		17.0	-0.365906194788025	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	MEHTHOFURAN	Shoot		85.0		Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	VERBENYL-ACETATE	Shoot		2.0	-0.8164965809277258	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-CARYOPHYLLENE	Shoot		32.0	-0.34509629975614076	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	GAMMA-MUUROLENE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALLO-AROMADENDRENE	Shoot		2.0	-0.18449899191288488	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BOURBONENE	Shoot		3.0	-1.1137070295367113	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	N-HEXANAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	EO	Shoot		1500.0	-0.8141965329475307	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	2,4-DECADIENAL	Shoot		2.0	1.0000000000000002	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALPHA-EUDESOL	Shoot		24.0	1.5508589875233674	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	CIS-BETA-OCIMENE	Shoot		8.0	-0.5039521391496569	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	1-OCTEN-3-OL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-FARNESENE	Shoot		16.0	-0.3715645380063965	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	LUTEOLIN-7-RUTINOSIDE	Plant				J.S. Glasby Dict. Pls Containing 2ndary Metabolite. 1991.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	VERBENYL-ACETATE	Shoot		3.0	1.2247448713915892	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-CARYOPHYLLENE	Shoot		75.0	-0.25792055501122796	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	GAMMA-MUUROLENE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALLO-AROMADENDRENE	Shoot		16.0	-0.1661040406381092	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BOURBONENE	Shoot		4.0	-0.7206339602884604	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	N-HEXANAL	Shoot		0.1	-1.0	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	(+)-ISOMENTHONE	Leaf	1230.0	3485.0	-0.08380547905273038	--

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	EO	Shoot		1100.0	-0.8512204032739488	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	2,4-DECADIENAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	TERPINENE	Leaf				List, P.H. and Horhammer, L., Hager's Handbuch der Pharmazeutischen Praxis, Vols. 2-6, Springer-Verlag, Berlin, 1969-1979.
0	ALPHA-EUDESOL	Shoot		22.0	1.3270236697364897	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	CIS-BETA-OCIMENE	Shoot		5.0	-0.5234943490109244	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	1-OCTEN-3-OL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-FARNESENE	Shoot		6.0	-0.4281479701901117	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	LUTEOLIN-7-BETA-D-GLUCOPYRANOSIDE	Plant				J.S. Glasby Dict.Plis Containing 2ndary Metabolite. 1991.
0	BETA-CYCLOCITRAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	TRANS-2-HEPTANAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-CARYOPHYLLENE	Shoot		115.0	-0.1768268389694486	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	GAMMA-MUUROLENE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALLO-AROMADENDRENE	Shoot		1.0	-0.18581291700394026	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BOURBONENE	Shoot		6.0	0.0655121782080416	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.



Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	MYRTENYL-METHYL-ETHER	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	EO	Leaf	3000.0	8500.0	-0.26696073465631487	List, P.H. and Horhammer, L., <i>Hager's Handbuch der Pharmazeutischen Praxis</i> , Vols. 2-6, Springer-Verlag, Berlin, 1969-1979.
0	2,4-DECADIENAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALPHA-EUDESOL	Shoot		7.0	-0.3517412136650936	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	CINEOLE	Leaf	63.0	1904.0	-0.5927137247457667	--
0	1-OCTEN-3-OL	Shoot		2.0	-0.6377271578856241	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	TRANS-2-HEPTANAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-CARYOPHYLLENE	Shoot		75.0	-0.25792055501122796	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	GAMMA-MUUROLENE	Shoot		1.0	-0.3724734981350398	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BICYCLOGERMACRENE	Shoot		2.0	-0.8626280704542847	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	MYRTENYL-METHYL-ETHER	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-CYCLOCITRAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	CIS-JASMONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	2,4-DECADIENAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	ALPHA-COPAENE	Shoot		2.0	-0.3247580887205161	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	1-OCTEN-3-OL	Shoot		0.1	-0.646273518203606	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	LINALOL	Leaf	1080.0	3060.0	1.3698944061795897	--
0	TRANS-2-HEPTANAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-CARYOPHYLLENE	Shoot		32.0	-0.34509629975614076	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ACACETIN-7-RUTINOSIDE	Plant				J.S. Glasby Dict.Pls Containing 2ndary Metabolite. 1991.
0	BICYCLOGERMACRENE	Shoot		5.0	-0.8497466159427722	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	MYRTENYL-METHYL-ETHER	Shoot		2.0	-0.9901799045609176	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	BETA-CYCLOCITRAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALPHA-GURJUNENE	Shoot		4.0	-0.9428785569529811	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	CIS-JASMONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	2,4-DECADIENAL	Shoot		0.1	-1.0	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALPHA-COPAENE	Shoot		2.0	-0.3247580887205161	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ISOAMYL-ISOVALERATE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	BETA-CYCLOCITRAL	Shoot		0.1	-0.7109289582578825	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	TRANS-2-HEPTANAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-CARYOPHYLLENE	Leaf	70.0	850.0	0.08134626903727397	--
0	BICYCLOGERMACRENE	Shoot		7.0	-0.841158979601764	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	MYRTENYL-METHYL-ETHER	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-CYCLOCITRAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALPHA-GURJUNENE	Shoot		8.0	-0.8502629278934796	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	CIS-JASMONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	OCTAN-3-OL-ACETATE	Leaf				Jim Duke's personal files.
0	1-OCTYL-3-YL-ACETATE	Shoot		2.0	-0.46090305337856907	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	(+)-NEOMENTHYL-ACETATE	Leaf	480.0	1360.0		--
0	ISOAMYL-ISOVALERATE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-CUBEBENE	Shoot		4.0	-0.253152042232558	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALPHA-COPAENE	Shoot		2.0	-0.3247580887205161	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	TRANS-2-HEPTANAL	Shoot		0.1		Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	APIGENIN-7-RUTINOSIDE	Plant				J.S. Glasby Dict.Pls Containing 2ndary Metabolite. 1991.
0	ERIODICTYOL-7-RUTINOSIDE	Plant				J.S. Glasby Dict.Pls Containing 2ndary Metabolite. 1991.
0	2-N-AMYL-FURAN	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
0	BICYCLOGERMACRENE	Shoot		6.0	-0.8454527977722681	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
0	MYRTENYL-METHYL-ETHER	Shoot		0.1	-1.0097721130343957	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
0	ALPHA-GURJUNENE	Shoot		12.0	-0.757647298833978	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
0	CIS-JASMONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of Mentha aquatica Containing Sesquiterpene Alcohols as Major Components. Nippon Nogeikagaku Kaishi 67(10): 1417-1419.
0	OCTAN-3-OL	Leaf	3.0	162.0	-0.4159560336666566	--

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	1-OCTYL-3-YL-ACETATE	Shoot		3.0	-0.43804835651682184	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	(+)-NEOMENTHOL	Leaf	630.0	1785.0	0.4755159366502996	--
0	ISOAMYL-ISOVALERATE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-CUBEBENE	Shoot		2.0	-0.2573198400961167	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALPHA-COPAENE	Shoot		2.0	-0.3247580887205161	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	APIGENIN-7-BETA-D-GLUCOPYRANOSIDE	Plant				J.S. Glasby Dict.Pls Containing 2ndary Metabolite. 1991.
0	ERIODICTYOL-7-BETA-D-GLUCOPYRANOSIDE	Plant				J.S. Glasby Dict.Pls Containing 2ndary Metabolite. 1991.
0	2-N-AMYL-FURAN	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	(+)-MENTHOFURAN	Leaf	330.0	1445.0	1.4129257812897391	--



Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	BICYCLOGERMACRENE	Shoot		2.0	-0.8626280704542847	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALPHA-GURJUNENE	Shoot		10.0	-0.8039551133637287	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	CIS-JASMONE	Shoot		0.1	-1.4988327508640107	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	1-OCTYL-3-YL-ACETATE	Shoot		2.0	-0.46090305337856907	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	N-HEXANAL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ISOAMYL-ISOVALERATE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	VERBENYL-ACETATE	Shoot		3.0	1.2247448713915892	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-CUBEBENE	Shoot		8.0	-0.2448164465054405	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALPHA-COPAENE	Shoot		2.0	-0.3247580887205161	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	EO	Shoot		1100.0	-0.8512204032739488	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	PIPERITONE-OXIDE	Leaf				Jim Duke's personal files.
0	2-N-AMYL-FURAN	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	(-)-LIMONENE	Leaf	210.0	595.0	-0.8265112371345806	--
0	ALPHA-GURJUNENE	Shoot		4.0	-0.9428785569529811	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	CIS-BETA-OCIMENE	Shoot		2.0	-0.5430365588721919	Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	1-OCTYL-3-YL-ACETATE	Shoot		3.0	-0.43804835651682184	Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BETA-FARNESENE	Shoot		6.0	-0.4281479701901117	Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	N-HEXANAL	Shoot				Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ISOAMYL-ISOVALERATE	Shoot		0.1	-0.4765294039940469	Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	VERBENYL-ACETATE	Shoot		2.0	-0.8164965809277258	Umamoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
0	BETA-CUBEBENE	Shoot		9.0	-0.2427325475736611	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	ALLO-AROMADENDRENE	Shoot		1.0	-0.18581291700394026	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	BOURBONENE	Shoot		6.0	0.0655121782080416	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	EO	Shoot		1600.0	-0.8049405653659262	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
0	2-N-AMYLURAN	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
1	VIRIDIFLOROL	Shoot		50.0	-0.24673262666016743	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
1	VIRIDIFLOROL	Shoot		140.0	1.109355092388081	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
1	VIRIDIFLOROL	Shoot		195.0	1.938075365139788	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
1	VIRIDIFLOROL	Shoot		135.0	1.0340168857742895	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
1	VIRIDIFLOROL	Shoot		140.0	1.109355092388081	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
1	VIRIDIFLOROL	Leaf	36.0	1105.0	4.397695972931276	--
1	MENTHYL-VALERATE	Leaf				ANON. 1948-1976. The Wealth of India raw materials. Publications and Information Directorate, CSIR, New Delhi. 11 volumes.
2	NONANOIC-ACID	Shoot		2.0	-0.4758133309661321	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	ELEMOL	Leaf	660.0	1870.0	1.3469294965542162	--

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
2	MENTHYL-ACETATE	Shoot		6.0	-0.49157346467694896	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	NONANOIC-ACID	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	MENTHYL-ACETATE	Shoot		2.0	-0.6750942248230098	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	ELEMOL	Shoot		70.0	-0.6732642434273933	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	NONANOIC-ACID	Shoot		0.1	-0.6146598831763856	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	GERMACRENE-D	Shoot		25.0	-0.35647627073349986	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
2	MENTHYL-ACETATE	Shoot		2.0	-0.6750942248230098	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	ELEMOL	Shoot		170.0	0.374889742188684	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	GERMACRENE-D	Shoot		80.0	-0.3256419647815558	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	MENTHYL-ACETATE	Shoot		6.0	-0.49157346467694896	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	ELEMOL	Shoot		275.0	1.4754514270855648	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	GERMACRENE-D	Shoot		130.0	-0.2976107775525158	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
2	BETA-MYRCENE	Shoot		12.0	-0.3360589101674067	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	MENTHYL-ACETATE	Leaf		1900.0	-1.0	--
2	ELEMOL	Shoot		220.0	0.8989667349967224	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	GERMACRENE-D	Shoot		115.0	-0.3060201337212278	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	BETA-MYRCENE	Shoot		26.0	-0.3074660526172515	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	ELEMOL	Shoot		70.0	-0.6732642434273933	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	GERMACRENE-D	Shoot		25.0	-0.35647627073349986	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.



Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
2	BETA-MYRCENE	Shoot		37.0	-0.2850002359707011	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	(-)-MENTHONE	Leaf	510.0	2805.0	-0.7890023610971657	--
2	BETA-MYRCENE	Shoot		36.0	-0.2870425829385694	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	NEOMENTHOL	Leaf				Jim Duke's personal files.
2	NONANOIC-ACID	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	BETA-MYRCENE	Shoot		85.0	-0.18696758151302673	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	NONANOIC-ACID	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
2	GERMACRENE-D	Leaf	70.0	595.0	-0.04345648866625705	--
3	ISOMENTHONE	Leaf	30.0	170.0	-0.62376353248996	--
3	ACETONE	Leaf				Jim Duke's personal files.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
3	ISOMENTHONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
3	ISOMENTHONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
3	ISOMENTHONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
3	METHYL-ACETATE	Shoot		16.0	1.0	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
3	ISOMENTHONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
3	PIPERITENONE	Leaf				Jim Duke's personal files.
3	ISOMENTHONE	Shoot		0.1	-0.5336018108879219	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
4	(+)-PULEGONE	Leaf		1445.0	-0.6679831237459577	--

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
5	BETA-ELEMENE	Shoot		65.0	-0.11953111910292297	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
5	BETA-ELEMENE	Shoot		65.0	-0.11953111910292297	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
5	BETA-ELEMENE	Shoot		29.0	-0.24164481962470882	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
5	SABINENE	Shoot		7.0	-0.2884688205415059	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
5	SABINENE	Shoot		35.0	-0.25865584465310376	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
5	SABINENE	Shoot		15.0	-0.2799508274305339	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
5	MENTHOFURAN	Shoot		500.0	-0.34612871699139214	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
5	SABINENE	Shoot		10.0	-0.2852745731248914	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
5	MENTHOFURAN	Shoot		395.0	-0.3669798445209941	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
5	SABINENE	Shoot		35.0	-0.25865584465310376	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
5	MENTHOFURAN	Shoot		180.0	-0.4096750104149409	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
5	MENTHOFURAN	Shoot		42.0	-0.4370793494538463	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
5	BETA-ELEMENE	Shoot		29.0	-0.24164481962470882	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
5	MENTHOFURAN	Leaf	30.0	7582.0		--
5	BETA-ELEMENE	Shoot		55.0	-0.1534515914700857	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
6	FURFURAL	Leaf				Jim Duke's personal files.
7	LINALYL-ACETATE	Leaf				List, P.H. and Horhammer, L., Hager's Handbuch der Pharmazeutischen Praxis, Vols. 2-6, Springer-Verlag, Berlin, 1969-1979.
7	PIPERITONE	Leaf				ANON. 1948-1976. The Wealth of India raw materials. Publications and Information Directorate, CSIR, New Delhi. 11 volumes.
8	HESPERETIN	Plant				J.S. Glasby Dict.Pls Containing 2ndary Metabolite. 1991.
8	CARYOPHYLLENE-OXIDE	Shoot		1.0	-0.3970730841170472	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
8	CARYOPHYLLENE-OXIDE	Shoot		3.0	-0.3926872831083181	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
8	CARYOPHYLLENE-OXIDE	Shoot		3.0	-0.3926872831083181	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
8	CARYOPHYLLENE-OXIDE	Shoot		6.0	-0.3861085815952244	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
8	CARYOPHYLLENE-OXIDE	Shoot		1.0	-0.3970730841170472	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
9	CAMPHENE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
9	TERPINOLENE	Shoot		2.0	-0.7480787235698596	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
9	CAMPHENE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
9	TERPINOLENE	Shoot		1.0	-0.7769620719702403	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
9	CAMPHENE	Shoot		0.1	-0.1342670938252942	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
9	TERPINOLENE	Shoot		1.0	-0.7769620719702403	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
9	CAMPHENE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
9	TERPINOLENE	Shoot		2.0	-0.7480787235698596	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
9	CAMPHENE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
9	TERPINOLENE	Shoot		2.0	-0.7480787235698596	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
11	NEROLIDOL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
11	NEROLIDOL	Shoot		3.0	-0.40119990016633905	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
11	NEROLIDOL	Shoot		2.0	-0.40423910148064757	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
11	NEROLIDOL	Shoot		4.0	-0.3981606988520306	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
11	NEROLIDOL	Shoot		0.1	-0.4100135839778336	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.



Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
11	GAMMA-TERPINENE	Shoot		2.0	-0.3929527157162362	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
11	GAMMA-TERPINENE	Shoot		5.0	-0.39087991166838365	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
11	GAMMA-TERPINENE	Shoot		5.0	-0.39087991166838365	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
11	GAMMA-TERPINENE	Shoot		2.0	-0.3929527157162362	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
11	GAMMA-TERPINENE	Shoot		8.0	-0.3888071076205311	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
13	BETA-PINENE	Shoot		27.0	-0.11920908974169492	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
13	BETA-PINENE	Shoot		20.0	-0.11969942581367655	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
13	BETA-PINENE	Shoot		13.0	-0.12018976188565819	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
13	BETA-PINENE	Leaf	15.0	315.0	0.035492687396899314	--
13	BETA-EUDES MOL	Shoot		8.0	-0.3290760279158904	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
13	BETA-EUDES MOL	Shoot		13.0	0.18510526570268904	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
13	BETA-EUDES MOL	Shoot		27.0	1.6248128878347115	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
13	FORMIC-ACID	Leaf				Jim Duke's personal files.
13	BETA-EUDES MOL	Shoot		24.0	1.3163041116635639	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
13	BETA-EUDESOL	Shoot		24.0	1.3163041116635639	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
13	BETA-PINENE	Shoot		13.0	-0.12018976188565819	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
13	BETA-PINENE	Shoot		24.0	-0.11941923377254418	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
16	P-CYMENE	Shoot		1.0	-0.500609925430423	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
16	P-CYMENE	Shoot		2.0	-0.4998171975740504	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
16	P-CYMENE	Shoot		2.0	-0.4998171975740504	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
16	P-CYMENE	Shoot		2.0	-0.4998171975740504	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
16	P-CYMENE	Shoot		1.0	-0.500609925430423	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
16	ACACETIN	Plant				J.S. Glasby Dict.Plis Containing 2ndary Metabolite. 1991.
19	CARVONE	Leaf	510.0	1785.0	0.08317670792891492	Jim Duke's personal files.
22	MYRCENE	Leaf				Jim Duke's personal files.
28	ALPHA-PINENE	Shoot		13.0	-0.14615199142269486	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
28	ALPHA-PINENE	Shoot		14.0	-0.14605225948006592	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
28	ALPHA-PINENE	Shoot		9.0	-0.1465509191932106	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
28	PULEGONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
28	ALPHA-PINENE	Shoot		8.0	-0.14665065113583953	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
28	PULEGONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
28	ALPHA-PINENE	Leaf	18.0	68.0	-0.22012965416485936	--
28	PULEGONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
28	PULEGONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
28	PULEGONE	Shoot		0.1	-0.2690227918468267	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
28	PULEGONE	Leaf	54.0	153.0	-0.6636583918928131	--
28	ALPHA-PINENE	Shoot		8.0	-0.14665065113583953	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
30	MENTHONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
30	MENTHONE	Shoot		2.0	-0.8211188672141561	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
30	MENTHONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
30	MENTHONE	Shoot		0.1	-0.8214418565250121	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
30	MENTHONE	Leaf	23.0	212.0	-0.613866095511492	--
30	MENTHONE	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
31	CARYOPHYLLENE	Leaf				Jim Duke's personal files.
35	TANNIN	Leaf		74000.0	-0.44596183266254885	List, P.H. and Horhammer, L., <i>Hager's Handbuch der Pharmazeutischen Praxis</i> , Vols. 2-6, Springer-Verlag, Berlin, 1969-1979.
53	LINALOOL	Leaf	1080.0	3060.0	1.471569724590626	List, P.H. and Horhammer, L., <i>Hager's Handbuch der Pharmazeutischen Praxis</i> , Vols. 2-6, Springer-Verlag, Berlin, 1969-1979.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
57	ROSMARINIC-ACID	Plant		27000.0	-0.08324448923511193	Fitoterapia No.62: 166.
60	LIMONENE	Shoot		60.0	-0.29789596175373306	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
60	LIMONENE	Shoot		75.0	-0.290260225754317	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
60	LIMONENE	Shoot		60.0	-0.29789596175373306	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
60	LIMONENE	Shoot		21.0	-0.31774887535221497	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
60	LIMONENE	Shoot		60.0	-0.29789596175373306	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
60	LIMONENE	Leaf	114.0	714.0	0.303344274117639	--
63	MENTHOL	Shoot		1.0	-0.6444587500209842	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.

Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
63	MENTHOL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
63	MENTHOL	Shoot		3.0	-0.6414348966484517	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
63	MENTHOL	Shoot				Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
63	MENTHOL	Shoot		1.0	-0.6444587500209842	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
63	MENTHOL	Leaf		3570.0	-0.19360438419646248	--
67	1,8-CINEOLE	Leaf	300.0	850.0	-0.42089267832815774	--
67	1,8-CINEOLE	Shoot		45.0	-0.13495308546531803	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
67	1,8-CINEOLE	Shoot		115.0	-0.13394916713007335	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.



Activities Count	Chemical	Plant Part	Low PPM	High PPM	StdDev	Reference Citation
67	1,8-CINEOLE	Shoot		140.0	-0.13359062486748594	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
67	1,8-CINEOLE	Shoot		110.0	-0.13402087558259082	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
67	1,8-CINEOLE	Shoot		110.0	-0.13402087558259082	Umemoto, K., Arai, T., Nii, H. and Furukawa, K. 1993. A New Chemotype of <i>Mentha aquatica</i> Containing Sesquiterpene Alcohols as Major Components. <i>Nippon Nogeikagaku Kaishi</i> 67(10): 1417-1419.
101	APIGENIN	Plant				J.S. Glasby <i>Dict.Pls Containing 2ndary Metabolite</i> . 1991.