

List of All Chemicals

Galactia jussiaeana (Fabaceae)

How used

Medicinal

*Unless otherwise noted all references are to Duke, James A. 1992. Handbook of phytochemical constituents of GRAS herbs and other economic plants. Boca Raton, FL. CRC Press.

Chemical	Part <input type="button" value="All"/>	Low PPM	High PPM	StdDev	*Reference
GENISTEIN	Leaf Diffusate	--	--		*
Activities (81)					
Abortifacient	Cooper-Driver, G. A., Chemical substances in plants toxic to animals, pp. 213-47 in Rechcigl, M., Jr., ed. CRC Handbook of Naturally Occurring Food Toxicants, CRC Press, Boca Raton, 1983, 339 pp.				
Aldose-Reductase-Inhibitor IC50=10 uM					
Alpha-Reductase-Inhibitor IC50=35 uM					
Antiaggregant 1-10 ug/ml	Medline (post 1990 searches filed in my computer)				
Antiangiogenic 150 uM					
Antiatherosclerotic	Journal of Medicinal Food 2: 163.1999.				
Anticancer (Breast) IC50=2.6-6.7 ug/ml					
Anticarcinomic (Breast) IC50=7640 uM	Medline (post 1990 searches filed in my computer)				
Anticlimacteric	Duke, J. A. Writeups or information summaries on approximately 2,000 economic plants, USDA, ARS, Beltsville, MD 20705.				
Antiendocytotic 100 ug/ml	Medline (post 1990 searches filed in my computer)				
Antiestrogenic					
Antifertility					
Antihemolytic					
Antiimplantation					
Antiinflammatory	Journal of Medicinal Food 2: 179.1999.				
Antiischemic	Medline (post 1990 searches filed in my computer)				
Antileukemic >50 ug/ml (185 uM)					
Antileukemic IC50=0.010 ug/ml	Life Sciences 55: 1061.				
Antileukotriene IC50=10-15 uM	Medline (post 1990 searches filed in my computer)				
Antilymphomic					
Antimelanomic	Medline (post 1990 searches filed in my computer)				
Antimicrobial					
Antimitogenic IC50=12 uM					
Antimutagenic ID50=50-100 nM	Maga, J.A. Contribution of Phenolic Compounds to Smoke Flavor. Phenolic Compounds in Food and Their Effects on Health, Ch.13.				
Antineuroblastomic					
Antiosteoporotic					
Antioxidant	Phenolic Compounds in Food and Their Effects on Health, 64.				
Antiproliferative 5-45 uM					
Antiprostataadenomic 8-27 ug/ml	Medline (post 1990 searches filed in my computer)				
Antiprostatic IC50=35 uM					
Antispasmodic IC50=34 uM					
Antitumor (Breast)					
Antitumor (GI)					
Antitumor (Ovary)					
Antitumor (Prostate)					
Antitumor (Stomach)					
Antiulcer?					
Antiviral					
Apoptotic					
Aromatase-Inhibitor IC30=1 uM/l	Journal of Medicinal Food 2: 235.1999.				
Calcium-Antagonist 50 uM gpg					
Cancer-Preventive	Stitt, P. A. Why George Should Eat Broccoli. Dougherty Co, Milwaukee, WI, 1990, 399 pp.				
Cardioprotective	Journal of Medicinal Food 2: 163.1999.				

Catechol-O-Methyltransferase-Inhibitor

Cytotoxic >50 ug/ml (185 uM) Medline (post 1990 searches filed in my computer)

Cytotoxic IC89=10 ug/ml Planta Medica, 57: A113, 1991.

DOPA-Decarboxylase-Inhibitor

Estrogen-Agonist 0.1-1 uM/l Journal of Medicinal Food 2: 139.1999.

Estrogenic EC50=0.1-25 uM/l

Flatulent

Fungicide 2.5 mM Geibel, M. 1994. Sensitivity of the Fungus *Cytospora personii* to the Flavonoids of *Prunus cerasus*. *Phytochemistry*. 38(3): 599-601. 1995.

Fungicide ED50=50->100

Fungistat IC34=200 uM

Histidine-Kinase-Inhibitor IC50=110 uM Medline (post 1990 searches filed in my computer)

Hypocholesterolemic ED50=0.5 mg/kg

Immunostimulant 2-20 mg/kg orl mus

Lipase-Inhibitor

MAO-Inhibitor IC50=60-140 uM

MAO-Inhibitor IC50=95

MDR-Inhibitor 200 uM Castro, A. F., Altenberg, G. A. 1997. Inhibition of Drug Transport by Genistein in Multidrug-Resistant Cell Expressing P-Glycoprotein. *Biochem Pharmacol*, 53(1): 89-93.

NADH-Ubiquinone-Oxidoreductase-Inhibitor

Natriuretic

Ornithine-Decarboxylase-Inhibitor

PTK-Inhibitor 10-100 uM

Peroxidase-Inhibitor Jeffery B. Harborne and H. Baxter, eds. 1983. *Phytochemical Dictionary. A Handbook of Bioactive Compounds from Plants*. Taylor & Frost, London. 791 pp.

Pesticide

Phytoalexin Nigg, H.N. and Seigler, D.S., eds. 1992. *Phytochemical Resources for Medicine and Agriculture*. Plenum Press, New York. 445 pp.

Pituitary-sensitizer Medline (post 1990 searches filed in my computer)

Quinone-Reductase-Inducer IC50=4.6 ug/ml

Topoisomerase-I-Inhibitor IC50=250 ppm

Topoisomerase-I-Inhibitor 1-10 ug/ml Okura, A., Arakawa, H., Oka, H., Yoshinari, T., Monden, Y. 1988. Effect of Genistein on Topoisomerase Activity and on the Growth of [VAL 12]HA-RAS-Transformed NIH 3T3 Cells. *Biochem Biophys Res Commun*, 157: 183-189.

Topoisomerase-II-Inhibitor IC50=1-150 uM

Topoisomerase-II-Inhibitor 2-7.5 ug/ml

Topoisomerase-II-Inhibitor IC50=30 ug/ml Constantinou, A., Mehta, R., Runyan, C., Rao, K., Vaughan, A., Moon, R. 1995. Flavonoids as DNA Topoisomerase Antagonists and Poisons: Structure-Activity Relationships. *J Natural Products*, 58: 217-225.

Topoisomerase-II-Inhibitor 20 uM Markovits, J., et al. 1989. Inhibitory Effects of the Tyrosine Kinase Inhibitor Genistein on Mammalian DNA Topoisomerase II. *Cancer Research*, 49: 5111-5117.

Topoisomerase-II-Poison Journal of Medicinal Food 2: 167.1999.

Trypanosomastat Medline (post 1990 searches filed in my computer)

Tyrosine-Kinase-Inhibitor

Uterotrophic ED50=0.5 mg/kg

Uterotrophic EC50=0.1-25 uM/l

VEGF-Inhibitor

HYDROXYGENISTEIN

Leaf Diffusate

--

--

*